

## 2.2 LIFE HISTORY INFORMATION AND LENGTH-DERIVED VARIABLES

The annual SAFE report will serve as the repository of available life history information for the Western Pacific region. Life history data, particularly age, growth, reproduction, and mortality information inform stock assessments on fish productivity and population dynamics. Some assessments, particularly for data poor stocks, utilize information from other areas that introduces biases and increases uncertainties in the population estimates. An archipelago-specific life history parameter ensures accuracy in the input parameters used in the assessment.

The NMFS PIFSC Bio-Sampling Program allows for the collection of life history samples like otoliths and gonads from priority species in the bottomfish and coral reef fisheries. A significant number of samples are also collected during research cruises. These life history samples, once processed and examined, will contribute to the body of scientific information for the two data-poor fisheries in the region (coral reef fish and bottomfish). The life history information available from the region will be monitored by the Fishery Ecosystem Plan Team and will be tracked through this section of the report.

This section will be divided into two fisheries: 1) prioritized coral reef ecosystem component species (ECS), and 2) management unit species (MUS). The prioritized coral reef species list was developed by the American Samoa Department of Marine and Wildlife Resources (DMWR) in 2019. The MUS are the species that are listed in the federal ecosystem plan and are managed on a federal level. Within each fishery, the available life history information will be described under the age, growth, and reproductive maturity section. The section labelled fish length derived parameters summarizes available information derived from sampling the fish catch or the market. Length-weight conversion coefficients provide area-specific values to convert length from fishery-dependent and fishery-independent data collection to weight or biomass.

### 2.2.1 American Samoa Coral Reef Ecosystem –Life History

#### 2.2.1.1 Age, Growth, and Reproductive Maturity

**Description:** Age determination is based on counts of yearly growth marks (annuli) and/or daily growth increments (DGIs) internally visible within transversely cut, thin sections of sagittal otoliths. Validated age determination is based on several methods including an environmental signal (bomb radiocarbon  $^{14}\text{C}$ ) produced during previous atmospheric thermonuclear testing in the Pacific and incorporated into the core regions of sagittal otolith and other aragonite-based calcified structures such as hermatypic corals. This technique relies on developing a regionally based aged coral core reference series for which the rise, peak, and decline of  $^{14}\text{C}$  values is available over the known age series of the coral core. Estimates of fish age are determined by projecting the  $^{14}\text{C}$  otolith core values back in time from its capture date to where it intersects with the known age  $^{14}\text{C}$  coral reference series. Fish growth is estimated by fitting the length-at-age data to a von Bertalanffy growth function. This function typically uses three coefficients ( $L_{\infty}$ ,  $k$ , and  $t_0$ ), which together characterize the shape of the length-at-age growth relationship.

Length-at-reproductive maturity is based on the histological analyses of small tissue samples of gonad material that are typically collected along with otoliths when a fish is processed for life history studies. The gonad tissue sample is preserved, cut into five-micron sections, stained, and sealed onto a glass slide for subsequent examination. Based on standard cell structure features and developmental stages within ovaries and testes, the gender, developmental stage, and maturity status (immature or mature) is determined via microscopic evaluation. The percent of

mature samples for a given length interval are assembled for each sex, and these data are fitted to a three- or four-parameter logistic function to determine the best fit for the data based on statistical analyses. The mid-point of the fitted function provides an estimate of the length at which 50% of fish have achieved reproductive maturity ( $L_{50}$ ). For species that undergo sex reversal (primarily female to male in the tropical Pacific region), such as groupers and deeper-water emperors among the bottomfishes, and for parrotfish, shallow-water emperors, and wrasses among the coral reef fishes, standard histological criteria are used to determine gender and reproductive developmental stages that indicate the transitioning or completed transition from one sex to another. These data are similarly analyzed using a three- or four-parameter logistic function to determine the best fit of the data based on statistical analyses. The mid-point of this fitted function provides an estimate of the length at which 50% of fish of a particular species have or are undergoing sex reversal ( $L\Delta_{50}$ ).

Age at 50% maturity ( $A_{50}$ ) and age at 50% sex reversal ( $A\Delta_{50}$ ) is typically derived by referencing the von Bertalanffy growth function for that species and using the corresponding  $L_{50}$  and  $L\Delta_{50}$  values to obtain the corresponding age value from this growth function. In studies where both age and growth and reproductive maturity are concurrently determined, estimates of  $A_{50}$  and  $A\Delta_{50}$  are derived directly by fitting the percent of mature samples for each age (i.e., one-year) interval to a three- or four-parameter logistic function using statistical analyses. The mid-point of this fitted logistic function provides a direct estimate of the age at which 50% of fish of a species have achieved reproductive maturity ( $A_{50}$ ) and sex reversal ( $A\Delta_{50}$ ).

**Data Category:** Biological

**Timeframe:** N/A

**Jurisdiction:** American Samoa

**Spatial Scale:** Archipelagic

**Data Source:** Sources of data are directly derived from research cruises sampling and market samples collected by the American Samoa contracted bio-sampling team which samples the catch of fishermen and local fish vendors. Laboratory analyses and data generated from these analyses reside with the PIFSC Life History Program. Refer to the “Reference” column in Table 18 for specific details on data sources by species.

**Parameter Definitions:**

**$T_{max}$  (maximum age)** – The maximum observed age revealed from an otolith-based age determination study.  $T_{max}$  values can be derived from ages determined by annuli counts of sagittal otolith sections and/or bomb radiocarbon ( $^{14}\text{C}$ ) analysis of otolith core material. Units are years.

**$L_{\infty}$  (asymptotic length)** – One of three coefficients of the von Bertalanffy growth function (VBGF) that measures the mean maximum length at which the growth curve plateaus and no longer increases in length with increasing age. This coefficient reflects the estimated mean maximum length and not the observed maximum length. Units are centimeters.

**$k$  (growth coefficient)** – One of three coefficients of the VBGF that measures the shape and steepness by which the initial portion of the growth function approaches its mean maximum length ( $L_{\infty}$ ).

**$t_0$  (hypothetical age at length zero)** – One of three coefficients of the VBGF whose measure is highly influenced by the other two VBGF coefficients ( $k$  and  $L_\infty$ ) and typically assumes a negative value when specimens representing early growth phases) are not available for age determination. This parameter can be fixed at 0. Units are years.

**$M$  (natural mortality)** – This is a measure of the mortality rate for a fish stock and is considered to be directly related to stock productivity (i.e., high  $M$  indicates high productivity and low  $M$  indicates low stock productivity).  $M$  can be derived through use of various equations that link  $M$  to  $T_{max}$  and the VBGF coefficients ( $k$  and  $L_\infty$ ) or by calculating the value of the slope from a regression fit to a declining catch curve (regression of the natural logarithm of abundance versus age class) derived from fishing an unfished or lightly fished population.

**$A_{50}$  (age at 50% maturity)** – Age at which 50% of the sampled stock under study has attained reproductive maturity. This parameter is best determined based on studies that concurrently determine both age (otolith-based age data) and reproductive maturity status (logistic function fitted to percent mature by age class with maturity determined via microscopic analyses of gonad histology preparations). A more approximate means of estimating  $A_{50}$  is to use an existing  $L_{50}$  estimate to find the corresponding age ( $A_{50}$ ) from an existing VBGF curve. Units are years.

**$A\Delta_{50}$  (age of sex switching)** – Age at which 50% of the immature and adult females of the sampled stock under study is undergoing or has attained sex reversal. This parameter is best determined based on studies that concurrently determines both age (otolith-based age data) and reproductive sex reversal status (logistic function fitted to percent sex reversal by age class with sex reversal determined via microscopic analyses of gonad histology preparations). A more approximate means of estimating  $A\Delta_{50}$  is to use an existing  $L\Delta_{50}$  estimate to find the corresponding age ( $A\Delta_{50}$ ) from the VBGF curve. Units are years.

**$L_{50}$  (length at which 50% of a fish population are capable of spawning)** – Length at which 50% of the females of a sampled stock under study has attained reproductive maturity; this is the length associated with  $A_{50}$  estimates. This parameter is derived using a logistic function to fit the percent mature data by length class with maturity status best determined via microscopic analyses of gonad histology preparations.  $L_{50}$  information is typically more available than  $A_{50}$  since  $L_{50}$  estimates do not require knowledge of age and growth. Units are centimeters.

**$L\Delta_{50}$  (length of sex switching)** – Length at which 50% of the immature and adult females of the sampled stock under study is undergoing or has attained sex reversal; this is the length associated with  $A\Delta_{50}$  estimates. This parameter is derived using a logistic function to fit the percent sex reversal data by length class with sex reversal status best determined via microscopic analyses of gonad histology preparations.  $L\Delta_{50}$  information is typically more available than  $A\Delta_{50}$  since  $L\Delta_{50}$  estimates do not require knowledge of age and growth. Units are centimeters.

**Rationale:** These nine life history parameters provide basic biological information at the species level to evaluate the productivity of a stock - an indication of the capacity of a stock to recover once it has been depleted. These parameters are also used as direct inputs into stock assessments. Currently, the assessment of coral reef fish resources in American Samoa is data-limited. Knowledge of these life history parameters support current efforts to characterize the resilience of these resources and provide important biological inputs for future stock assessment efforts and enhance our understanding of the species' likely role and status as a component of the overall ecosystem. Furthermore, knowledge of life histories across species at the taxonomic level of families or among different species that are ecologically or functionally similar can provide

important information on the diversity of life histories and the extent to which species can be grouped (based on similar life histories) for future multi-species assessments.

**Table 1. Available age, growth, and reproductive maturity information for prioritized coral reef ecosystem species in American Samoa**

Species	Age, growth, reproductive maturity parameters									Reference
	$T_{max}$	$L_{\infty}$	$k$	$t_0$	$M$	$A_{50}$	$A\Delta_{50}$	$L_{50}$	$L\Delta_{50}$	
<i>Cernimugil crenilabis</i>										
<i>Epinephelus melanostigma</i>										
<i>Octopus cyanea</i>										
<i>Panulirus peniciliatus</i>										
<i>Sargocentron tiere</i>										
<i>Tridacna maxima</i>										

<sup>a</sup> signifies estimate pending further evaluation in an initiated and ongoing study.

<sup>b</sup> signifies a preliminary estimate taken from ongoing analyses.

<sup>c</sup> signifies an estimate documented in an unpublished report or draft manuscript.

<sup>d</sup> signifies an estimate documented in a finalized report or published journal article (including in press).

Parameter estimates are for females unless otherwise noted (F=females, M=males). Parameters  $T_{max}$ ,  $t_0$ ,  $A_{50}$ , and  $A\Delta_{50}$  are in years;  $L_{\infty}$ ,  $L_{50}$ , and  $L\Delta_{50}$  are in mm fork length (FL);  $k$  is in units of year<sup>-1</sup>; X means the parameter estimate is too preliminary and Y means the published age and growth parameter estimates are based on DGI numerical integration technique and likely to be inaccurate. Superscript letters indicate status of parameter estimate (see footnotes below table). Published or in press publications (<sup>d</sup>) are shown in “Reference” column.

### 2.2.1.2 Fish Length Derived Parameters

**Description:** The NMFS Commercial Fishery Bio-Sampling Program started in 2010 and ended in 2015. This program had two components: first was the Field/Market Sampling Program and the second was the Lab Sampling Program, details of which are described in a separate section of this report. The goals of the Field/Market Sampling Program were to:

- Broad scale looks at commercial landings (by fisher/trip, gear, and area fished);
- Length and weight frequencies of whole commercial landings per fisher-trip (with an effort to also sample landings not sold commercially);
- Accurate species identification; and
- Develop accurate local length-weight curves.

In American Samoa, the Bio-Sampling Program focused on the commercial coral reef spear fishery with occasional sampling of the bottomfish fishery occurring locally and less frequently at the northern islands. Sampling is conducted in partnership with the fish vendors. The Market

Sampling information includes (but not limited to): 1) fish length; 2) fish weight; 3) species identification; and 4) basic effort information.

**Data Category:** Biological

**Timeframe:** N/A

**Jurisdiction:** American Samoa

**Spatial Scale:** Archipelagic

**Data Source:** NMFS Bio-Sampling Program

**Parameter Definitions:**

*n* – **sample size** is the total number of samples accumulated for each species recorded in the Bio-Sampling Program database from the commercial spear fishery.

*L<sub>max</sub>* – **maximum fish length** is the largest individual per species recorded in the Bio-Sampling Program database from the commercial spear fishery. This value is derived from measuring the length of individual samples for species occurring in the spear fishery. Units are centimeters.

*N<sub>L-W</sub>* – **sample size for L-W regression** is the number of samples used to generate the *a* and *b* coefficients.

*a* and *b* – **length-weight coefficients** are the coefficients derived from the regression line fitted to all length and weight measured by species in the commercial spear fishery. These values are used to convert length information to weight. Values are influenced by the life history characteristics of the species, geographic location, population status, and nature of the fisheries from which the species are harvested.

**Rationale:** Length-derived information is an important component of fisheries monitoring and data poor stock assessment approaches. Maximum length (*L<sub>max</sub>*), is used to derive missing species- and location-specific life history information (Nadon et al. 2015, Nadon and Ault 2016, Nadon 2019). The length-weight coefficients (*a* and *b* values) are used to convert length to weight for fishery-dependent and fishery-independent data collection where length is typically recorded but weight is the factor being used for management. This section of the report presents the best available information for the length-derived variables for the American Samoa coral reef fisheries.

**Table 2. Available length-derived information for prioritized coral reef species in American Samoa.**

Species	Length-derived parameters					Reference
	<i>n</i>	<i>L<sub>max</sub></i>	<i>N<sub>L-W</sub></i>	<i>a</i>	<i>b</i>	
<i>Crenimugil crenilabis</i>	380	48.2	380	0.0388	2.73	Matthews et al. 2019
<i>Epinephelus melanostigma</i>	2,662	54.9	2,662	0.0109	3.10	Matthews et al. 2019
<i>Octopus cyanea</i>						
<i>Panulirus peniciliatus</i>	3,384	15.8	3,384	2.6004	2.41	Matthews et al. 2019
<i>Sargocentron tiere</i>	3,002	25.0	3,002	0.069	2.62	Matthews et al. 2019
<i>Tridacna maxima</i>						

## 2.2.2 American Samoa Ecosystem – Management Unit Species Life History

### 2.2.2.1 Age, Growth, and reproductive Maturity

**Description:** Age determination is based on counts of yearly growth marks (annuli) and/or daily growth increments (DGIs) internally visible within transversely cut, thin sections of sagittal otoliths. Validated age determination is based on several methods including an environmental signal (bomb radiocarbon  $^{14}\text{C}$ ) produced during previous atmospheric thermonuclear testing in the Pacific and incorporated into the core regions of sagittal otolith and other aragonite-based calcified structures such as hermatypic corals. This technique relies on developing a regionally based aged coral core reference series for which the rise, peak, and decline of  $^{14}\text{C}$  values is available over the known age series of the coral core. Estimates of fish age are determined by projecting the  $^{14}\text{C}$  otolith core values back in time from its capture date to where it intersects with the known age  $^{14}\text{C}$  coral reference series. Fish growth is estimated by fitting the length-at-age data to a von Bertalanffy growth function. This function typically uses three coefficients ( $L_{\infty}$ ,  $k$ , and  $t_0$ ), which together characterize the shape of the length-at-age growth relationship.

Length-at-reproductive maturity is based on the histological analyses of small tissue samples of gonad material that are typically collected along with otoliths when a fish is processed for life history studies. The gonad tissue sample is preserved, cut into five-micron sections, stained, and sealed onto a glass slide for subsequent examination. Based on standard cell structure features and developmental stages within ovaries and testes, the gender, developmental stage, and maturity status (immature or mature) is determined via microscopic evaluation. The percent of mature samples for a given length interval are assembled for each sex, and these data are fitted to a three- or four-parameter logistic function to determine the best fit for the data based on statistical analyses. The mid-point of the fitted function provides an estimate of the length at which 50% of fish have achieved reproductive maturity ( $L_{50}$ ). For species that undergo sex reversal (primarily female to male in the tropical Pacific region), such as groupers and deeper-water emperors among the bottomfishes, and for parrotfish, shallow-water emperors, and wrasses among the coral reef fishes, standard histological criteria are used to determine gender and reproductive developmental stages that indicate the transitioning or completed transition from one sex to another. These data are similarly analyzed using a three- or four-parameter logistic function to determine the best fit of the data based on statistical analyses. The mid-point of this fitted function provides an estimate of the length at which 50% of fish of a particular species have or are undergoing sex reversal ( $L\Delta_{50}$ ).

Age at 50% maturity ( $A_{50}$ ) and age at 50% sex reversal ( $A\Delta_{50}$ ) is typically derived by referencing the von Bertalanffy growth function for that species and using the corresponding  $L_{50}$  and  $L\Delta_{50}$  values to obtain the corresponding age value from this growth function. In studies where both age and growth and reproductive maturity are concurrently determined, estimates of  $A_{50}$  and  $A\Delta_{50}$  are derived directly by fitting the percent of mature samples for each age (i.e., one-year) interval to a three- or four-parameter logistic function using statistical analyses. The mid-point of this fitted logistic function provides a direct estimate of the age at which 50% of fish of a species have achieved reproductive maturity ( $A_{50}$ ) and sex reversal ( $A\Delta_{50}$ ).

**Category:** Biological

**Timeframe:** N/A

**Jurisdiction:** American Samoa

**Spatial Scale:** Archipelagic

**Data Source:** Sources of data are directly derived from field samples collected at sea on NOAA research vessels and from the American Samoa contracted bio-sampling team which samples the catch of fishermen and local fish vendors. Laboratory analyses and data generated from these analyses reside with the PIFSC Life History Program. Refer to the “Reference” column in Table 21 for specific details on data sources by species.

**Parameter definitions:** Identical to Section 2.2.1.1.

**Rationale:** These nine life history parameters provide basic biological information at the species level to evaluate the productivity of a stock - an indication of the capacity of a stock to recover once it has been depleted. Currently, the assessment of coral reef fish resources in American Samoa is data-limited. Knowledge of these life history parameters support current efforts to characterize the resilience of these resources and provide important biological inputs for future stock assessment efforts and enhance our understanding of the species likely role and status as a component of the overall ecosystem. Furthermore, knowledge of life histories across species at the taxonomic level of families or among different species that are ecologically or functionally similar can provide important information on the diversity of life histories and the extent to which species can be grouped (based on similar life histories) for future multi-species assessments.

**Table 3. Available age, growth, and reproductive maturity information for MUS targeted for otoliths and gonads sampling in American Samoa**

Species	Age, growth, and reproductive maturity parameters									Reference
	$T_{max}$	$L_{\infty}$	$k$	$t_0$	$M$	$A_{50}$	$L_{50}$	$L\Delta_{50}$		
<i>Aphareus rutilans</i>	X <sup>a</sup>	X <sup>a</sup>	X <sup>a</sup>	X <sup>a</sup>	X <sup>a</sup>		NA		NA	LHP (in prep)
<i>Aprion virescens</i>							NA		NA	
<i>Caranx lugubris</i>							NA		NA	
<i>Etelis carbunculus</i> <sup>1</sup>							NA		NA	
<i>Etelis coruscans</i>							NA		NA	
<i>Lethrinus rubrioperculatus</i>										
<i>Lutjanus kasmira</i>							NA		NA	
<i>Pristipomoides filamentosus</i>							NA		NA	
<i>Pristipomoides flavipinnis</i>	28	41.15	0.47		0.22		NA		NA	O'Malley et al. 2019
<i>Pristipomoides zonatus</i>							NA		NA	
<i>Variola louti</i>										

<sup>1</sup> *E. carbunculus* is now known to be comprised of two distinct, non-interbreeding lineages (Andrews et al. 2016). Both species occur in the Samoa Archipelago and were likely both captured by fishermen in the 1980s but reported as one species.

<sup>a</sup> signifies estimate pending further evaluation in an initiated and ongoing study.

<sup>b</sup> signifies a preliminary estimate taken from ongoing analyses.

<sup>c</sup> signifies an estimate documented in an unpublished report or draft manuscript.

<sup>d</sup> signifies an estimate documented in a finalized report or published journal article (including in press).

### 2.2.2.2 Fish Length Derived Parameters

**Description:** The NMFS Commercial Fishery Bio-Sampling Program started in 2010 and ended in 2015. This program had two components: first was the Field/Market Sampling Program and the second was the Lab Sampling Program, details of which are described in a separate section of this report. The goals of the Field/Market Sampling Program were:

- Broad scale looks at commercial landings (by fisher/trip, gear, and area fished);
- Length and weight frequencies of whole commercial landings per fisher-trip (with an effort to also sample landings not sold commercially);
- Accurate species identification; and
- Develop accurate local length-weight curves.

In American Samoa, the Bio-Sampling focused on the commercial coral reef spear fishery with occasional sampling of the bottomfish fishery occurring locally and less frequently at the northern islands. Sampling was conducted in partnership with the fish vendors. The Market Sampling information includes (but not limited to): 1) fish length; 2) fish weight; 3) species identification; and 4) basic effort information.

**Category:** Biological

**Timeframe:** N/A

**Jurisdiction:** American Samoa

**Spatial Scale:** Archipelagic

**Data Source:** NMFS BioSampling Program

**Parameter Definition:** Identical to Section 2.2.1.2.

**Rationale:** Length-derived information is an important component of fisheries monitoring and data poor stock assessment approaches. Maximum length ( $L_{max}$ ), is used to derive missing species- and location-specific life history information (Nadon et al. 2015, Nadon and Ault 2016, Nadon 2019). The length-weight coefficients ( $a$  and  $b$  values) are used to convert length to weight for fishery-dependent and fishery-independent data collection where length is typically recorded but weight is the factor being used for management. This section of the report presents the best available information for the length-derived variables for the American Samoa MUS fishery.

**Table 4. Available length-derived information for MUS in American Samoa**

Species	Length derived parameters					Reference
	$n$	$L_{max}$	$N_{L-w}$	$a$	$b$	
<i>Aphareus rutilans</i>	173	85.0	173	0.0395	2.73	Matthews et al. 2019
<i>Aprion virescens</i>	952	73.4	952	0.0157	2.99	Matthews et al. 2019
<i>Caranx lugubris</i>	164	86.0	164	0.0404	2.80	Matthews et al. 2019
<i>Etelis carbunculus</i> <sup>1</sup>						
<i>Etelis coruscans</i>	106	89.5	106	0.0322	2.81	Matthews et al. 2019
<i>Lethrinus rubrioperculatus</i>	2,349	57.0	2,349	0.0287	2.86	Matthews et al. 2019
<i>Lutjanus kasmira</i>	461	35.0	461	0.0176	3.01	Matthews et al. 2019
<i>Pristipomoides flavipinnis</i>	262	56.5	262	0.0249	2.90	Matthews et al. 2019



<i>Pristipomoides zonatus</i>						
<i>Pristipomoides filamentosus</i>						
<i>Variola louti</i>		50.5	365	0.0135	3.08	Matthews et al. 2019

<sup>1</sup> *E. carbunculus* is now known to be comprised of two distinct, non-interbreeding lineages (Andrews et al. 2016). Both species occur in the Samoa Archipelago and were likely both captured by fishermen in the 1980s but reported as one species.

## Literature Cited

Matthews, T., D. Ochavillo, S. Felise, T. Letalie, M. Letuane, E. Schuster, A. Soonaolo, S. Tofaeono, A. Tua, and F. Tuilagi. 2019. Length-weight relationships for 71 reef and bottomfish species from Tutuila and Aunu'u, American Samoa. Pacific Islands Fish. Sci. Cent., Natl. Mar. Fish. Serv., NOAA, Honolulu, HI 96818-5007. Pacific Islands Fish. Sci. Cent. Admin. Rep. H-19-03, 9 p.

Nadon, M.O., Ault, J.S., Williams, I.D., Smith, S.G., and G.T. DiNardo. 2015. Length-based assessment of coral reef fish populations in the Main and Northwestern Hawaiian Islands. PLoS One, 10(8), p.e0133960.

Nadon MO, Ault JS. 2016. A stepwise stochastic simulation approach to estimate life history parameters for data-poor fisheries. Canadian Journal of Fisheries and Aquatic Sciences. <https://doi.org/10.1139/cjfas-2015-0303>.

Nadon, MO. 2019. Stock Assessment of Guam Coral Reef Fish, 2019. NOAA Technical Memorandum NMFS-PIFSC-82, 107 p. doi: 10.25923/pyd6-7k49.

O'Malley, J.M., C.B. Wakefield, Z. Oyafuso, R.S. Nichols, B.M. Taylor, A. J. Williams, M. Sapatu, M. Marsik. Effects of exploitation evident in age-based demography of 2 deepwater snappers, the goldeneye jobfish (*Pristipomoides flavipinnis*) in the Samoa Archipelago and the goldflag jobfish (*P. auricilla*) in the Mariana Archipelago. Fisheries Bulletin 117:322–336 doi: 10.7755/FB.117.4.5.