

**ANNUAL STOCK ASSESSMENT AND FISHERY
EVALUATION REPORT:
PACIFIC REMOTE ISLAND AREA
FISHERY ECOSYSTEM PLAN
2018**



Western Pacific Regional Fishery Management Council
1164 Bishop St., Suite 1400
Honolulu, HI 96813

PHONE: (808) 522-8220
FAX: (808) 522-8226
www.wpcouncil.org

The ANNUAL STOCK ASSESSMENT AND FISHERY EVALUATION REPORT for the PACIFIC REMOTE ISLAND AREA FISHERY ECOSYSTEM PLAN 2018 was drafted by the Fishery Ecosystem Plan Team. This is a collaborative effort primarily between the Western Pacific Regional Fishery Management Council (WPRFMC), National Marine Fisheries Service (NMFS)-Pacific Island Fisheries Science Center (PIFSC), Pacific Islands Regional Office (PIRO), Division of Aquatic Resources (HI) Department of Marine and Wildlife Resources (American Samoa), Division of Aquatic and Wildlife Resources (Guam), and Division of Fish and Wildlife (CNMI).

This report attempts to summarize annual fishery performance looking at trends in catch, effort and catch rates as well as provide a source document describing various projects and activities being undertaken on a local and federal level. The report also describes several ecosystem considerations including fish biomass estimates, biological indicators, protected species, habitat, climate change, and human dimensions. Information like marine spatial planning and best scientific information available for each fishery are described. This report provides a summary of annual catches relative to the Annual Catch Limits established by the Council in collaboration with the local fishery management agencies.

Edited By: Thomas Remington, Marlowe Sabater, Asuka Ishizaki, and Sylvia Spalding, WPRFMC.

This document can be cited as follows:

WPRFMC, 2019. Annual Stock Assessment and Fishery Evaluation Report: Pacific Remote Island Areas Fishery Ecosystem Plan 2018. Sabater, M., Ishizaki, A., Remington, T., and Spalding, S. (Eds.) Western Pacific Regional Fishery Management Council. Honolulu, Hawaii 96813 USA. 92 pp. + Appendices.

The **Western Pacific Regional Fishery Management Council** acknowledges the valuable contributions of the following Plan Team members for drafting sections of this report:

NMFS Pacific Islands Fisheries Science Center: Justin Hospital, Ivor Williams, Joe O'Malley, Michael Parke, Tom Oliver, Hannah Barkley, Frank Parrish, T. Todd Jones, Minling Pan, and Kirsten Leong.

NMFS Pacific Islands Regional Office: Brett Schumacher and Rebecca Walker.

The Council also acknowledges the staff of the **NMFS PIFSC Western Pacific Fisheries Information Network (WPacFIN)** for providing the technical support to generate the data summaries.

The Council would like to thank the following individuals for their contributions to the report: Samuel Kahng and Alexander Filardo.

This page was intentionally left blank.

EXECUTIVE SUMMARY

As part of its five-year fishery ecosystem plan (FEP) review, the Council identified the annual reports as a priority for improvement. The former annual reports have been revised to meet National Standard regulatory requirements for the Stock Assessment and Fishery Evaluation (SAFE) reports. The purpose of the reports is twofold: to monitor the performance of the fishery and ecosystem to assess the effectiveness of the FEP in meeting its management objectives, and to maintain the structure of the FEP living document. The reports are typically comprised of three chapters: fishery performance, ecosystem considerations, and data integration. The Council will iteratively improve the Annual SAFE Report as resources allow. The 2018 Pacific Remote Island Area (PRIA) FEP Annual SAFE Report does not contain fully-developed fishery performance or data integration chapters due to the absence of consistent fisheries data in the PRIAs.

An ecosystem considerations section was added to the annual SAFE report following the Council's review of its FEPs and revised management objectives (pending Secretarial transmittal). Fishery independent ecosystem survey, socioeconomic, protected species, oceanic and climate indicator, essential fish habitat, and marine planning information are all included in the ecosystem considerations section. Fishery dependent data sections will continue to be included and updated as resources allow.

Fishery independent ecosystem data were acquired through visual surveys conducted in the PRIAs, American Samoa, Guam, the Commonwealth of the Northern Mariana Islands (CNMI), the Main Hawaiian Islands (MHI), and the Northwestern Hawaiian Islands (NWHI). This report describes mean fish biomass for coral reefs in each of these locations. Additionally, the mean reef fish biomass and mean size of fishes (> 10 cm) in the PRIAs are presented by sampling year and reef area. Finally, the reef fish population estimates across hard bottom habitat (0-30 m) are provided for each PRIA study site.

The socioeconomics section is meant to outline the pertinent economic, social, and community information available for assessing the successes and impacts of management measures or the achievements of the FEP within the PRIAs. The section begins with an overview of the socioeconomic context for the region, and then provides a summary of relevant studies for the PRIAs. Because human habitation is limited in the PRIAs, socioeconomic information is also limited. The socioeconomics section of this report will be expanded in later years if activity increases and as resources allow. There were no new socioeconomic data reported from the PRIAs in 2018.

The protected species section of this report describes monitoring and summarizes protected species interactions in fisheries managed under the PRIA FEP. There are currently no bottomfish, crustacean, or precious coral fisheries operating in the PRIAs and no historical observer data are available for fisheries under this FEP. No new fishing activity was reported in 2018, and there is no other information to indicate that impacts to protected species from PRIA fisheries have changed over the past year.

The climate change section of this report includes measurements of changing climatic and related oceanic conditions in the geographic areas that the Western Pacific Regional Fishery Management Council has jurisdiction. In developing this section, the Council relied on a number of recent reports conducted in the context of the U.S. National Climate Assessment including the 2012 Pacific Islands Regional Climate Assessment as well as the ‘Ocean and Coasts’ chapter of the 2014 Pilot Indicator Systems report prepared by the National Climate Assessment and Development Advisory Committee. The primary goal for selecting the climatic indicators used in this report is to provide fisheries-related communities, resource managers, and businesses with climate-related situational awareness. In this context, indicators were selected to be fishery-relevant, be informative, build intuition about current conditions in light of changing climate, provide historical context, and distinguish patterns and trends.

The trend of atmospheric concentration of carbon dioxide (CO₂), for example, is increasing exponentially with a time series maximum at 409 ppm. Since 1989, the oceanic pH at Station Aloha in Hawaii has shown a significant linear decrease of -0.0389 pH units, or roughly a 9.4% increase in acidity ([H⁺]). The Oceanic Niño Index, which is a measure of the El Niño – Southern Oscillation (ENSO) phase, was mostly neutral in 2018, but the Pacific Decadal Oscillation (PDO) was positive (i.e., warm) for most of the year. The Accumulated Cyclone Energy (ACE) Index (x 10⁴ kt²) was well above the 30-year mean in both the Central and Eastern North Pacific. The Central North Pacific had six named storms where all six became hurricanes and three became major, while the Eastern North Pacific had 23 named storms where 13 became hurricanes and 10 became major hurricanes. The mean sea surface temperature (SST) data from the PRIAs showed annual anomalies that were slightly hotter than average. After a major heat stress events in 2015 and 2016 in all three virtual stations, the Northern Line Island virtual station showed a minor coral thermal stress exposure event in 2018 that reached nearly five degree heating weeks (DHW). The chlorophyll-a concentrations around the PRIAs were approximately in line with previous climatological ranges, while the precipitation anomalies were slightly positive. Sea level rise is roughly 0.75 mm/year on Johnston Atoll and 2.1 mm/year on Wake Island.

The effective fish habitat (EFH) section of the 2018 Annual SAFE Report for the PRIA FEP includes responses to previous Council recommendations regarding EFH, habitat use by MUS in the PRIAs, trends in habitat conditions, and cumulative impacts on EFH. Guidelines also require a report on the condition of the habitat; mapping progress and benthic cover are included as preliminary indicators pending development of habitat condition indicators for the PRIAs not otherwise represented in other sections of this report. The mean percent cover of live coral, macroalgae, and crustose coralline algae from RAMP sites collected from towed-diver surveys in the PRIA are also presented for the available years between 2001 and 2016. Levels of available EFH information are summarized for bottomfish, crustacean, and precious coral MUS. An EFH review of CREMUS has not been undertaken, as the Council recently completed its process incorporating certain CREMUS into the ecosystem component species (ECS) classification that does not require EFH designations. The Annual SAFE Report also addresses any Council directives toward its Plan Team, though there were no directives in 2018 for the PRIAs.

The marine planning section of the 2018 annual SAFE report tracks activities with multi-year planning horizons and begins to monitor the cumulative impact of established facilities. The

Pacific Islands Marine National Monument (PIMNM) remains intact around the islands and atolls of the PRIAs. In June 2018, President Trump signed the EO 13840 *Regarding the Ocean Policy to Advance Economic, Security, and Environmental Interests of the United States*, which revoked EO 13547, eliminated the mandate for the federal government to participate in ocean planning at a regional level, and disbanded the associated regional planning bodies (RPBs). EO 13840 also requires federal agencies to coordinate activities regarding ocean-related matters and facilitate the coordination and collaboration of ocean-related matters with governments and ocean stakeholders, prompting the creation of the American Samoa Coastal and Marine Spatial Planning Data Portal by Marine Cadastre. The portal is intended to be expanded to incorporate the rest of the Western Pacific region, including the PRIAs, and be titled the Pacific Islands Regional Marine Planner. Development of the report in later years will focus on identifying appropriate data streams to be presented. No new ocean activities with multi-year planning horizons were identified for the PRIA in 2018.

The data integration chapter of this report is not fully developed. Previously, in late 2016, the Council hosted a data integration workshop with participants from the NMFS PIRO and PIFSC to identify policy-relevant fishery ecosystem relationships. However, no updates were made for the PRIA data integration chapter for the 2018 Annual SAFE Report. Despite the presence of data for certain ecological parameters throughout the remote Pacific Islands, there exists no fishery performance data in the absence of active fisheries and associated fishery-dependent information streams. The data integration chapter will be expanded in later years if fishing activity and data availability increases in the PRIAs.

Regarding the revisions to the 2018 Annual FEP SAFE Reports, the 2019 Archipelagic Plan Team recommends the Council:

- Direct staff to work with NMFS to convene the Plan Team working group for American Samoa, Guam, CNMI, and Hawaii to define the ecosystem component species that will be monitored as species that comprise the functional groups (e.g., ‘parrotfish’, ‘browsing surgeon’, ‘mid-size targeted surgeon’, ‘medium large snappers’, ‘non-planktivorous butterflyfishes’), and those that comprise key species in the fisheries (i.e., top 5 consistently monitored important species and the 10 annual catch landings)
- Direct staff to work with NMFS and American Samoa Department Marine Wildlife Resources (AS-DMWR), CNMI Division of Fish and Wildlife (DFW), Guam Division of Aquatic and Wildlife Resources (DAWR), Hawaii-DAR on the revisions to the fisheries modules of the Archipelagic SAFE Reports due to the changes in the Management Unit Species brought about by the Ecosystem Component designation; and
- Direct staff to work with NMFS-PIFSC-Ecosystem Science Division and Division of Aquatic Resources on applying the general linear modelling (GLM) framework to the survey data in order to validate the modeling results.

Additional work item recommendations include:

- Council staff and the Archipelagic Plan Team Chair to work with NMFS and AS-DMWR, CNMI-DFW, Guam-DAWR, Hawaii-DAR on determining which table(s) to remove from the Annual SAFE Report due to the ecosystem component amendment, etc.;
- WPacFIN to follow-up on the status of the creel survey method documentation;
- The report to incorporate more nuance in the narratives of the fishery performance sections; include the issue on pounds sold greater than pounds caught;
- The report to identify presence and absence of hi-liners in the data sets as well as define the criteria of what a hi-liner is;
- Regarding effort and participation metrics for the Annual SAFE Report, Council staff and PIFSC employees to calculate the average fishermen per trip and ensure interview has number of fishermen and average numbers of gear per trip.
- Add the abstracts for relevant data integration studies to Chapter 3; and
- “Cross-walk” tables with the information regularly needed to complete Environmental Assessments (EAs).

TABLE OF CONTENT

Acronyms and Abbreviations	xiv
1 Fishery Performance	1
1.1 Number of Federal Permit Holders.....	1
1.1.1 Special Coral Reef Ecosystem Permit	1
1.1.2 Western Pacific Precious Corals Permit	1
1.1.3 Western Pacific Crustaceans Permit (Lobster or Deepwater Shrimp).....	1
1.1.4 PRIA Bottomfish Permit.....	1
1.2 Administrative and Regulatory Actions	2
2 Ecosystem Considerations	3
2.1 Coral Reef Fish Ecosystem Parameters	3
2.1.1 Regional Reef Fish Biomass.....	3
2.1.2 Archipelagic Reef Fish Biomass.....	6
2.1.3 Archipelagic Mean Fish Size.....	9
2.1.4 Reef Fish Population Estimates	24
2.1.5 References.....	25
2.2 Protected Species	26
2.2.1 Monitoring Protected Species Interactions in the PRIA FEP Fisheries.....	26
2.2.2 Status of Protected Species Interactions in the PRIA FEP Fisheries.....	28
2.2.3 Identification of Emerging Issues	28
2.2.4 Identification of Research, Data, and Assessment Needs.....	30
2.3 Socioeconomics	31
2.3.1 Response to Previous Council Recommendations.....	32
2.3.2 Background.....	32
2.3.3 Ongoing Research and Information Collection	32
2.3.4 Relevant PIFSC Economics and Human Dimensions Publications: 2018	32
2.3.5 References.....	33
2.4 Climate and Oceanic Indicators.....	34
2.4.1 Introduction.....	34
2.4.2 Conceptual Model.....	35
2.4.3 Selected Indicators	37
2.5 Essential Fish Habitat	69
2.5.1 Introduction.....	69
2.5.2 EFH Information.....	69
2.5.3 Habitat Use by MUS and Trends in Habitat Condition	71
2.5.4 Report on Review of EFH Information	79
2.5.5 EFH Levels	79
2.5.6 Research and Information Needs.....	82
2.5.7 References.....	83

2.6	Marine Planning.....	85
2.6.1	Introduction.....	85
2.6.2	Activities and Facilities	89
2.6.3	Pacific Islands Regional Planning Body Report.....	89
2.6.4	References.....	89
3	Data Integration	90
Appendix A: PRIA FEP management unit species list.....		A-1
Appendix B: List of Protected Species and Designated Critical Habitat.....		B-1

TABLE OF TABLES

Table 1. Number of federal permit holders in the lobster, shrimp, and bottomfish fisheries of the PRIA from 2009 to 2018..... 2

Table 2. Reef fish population estimates for PRIA CREMUS in 0-30 m hard bottom habitat..... 24

Table 3. Summary of ESA consultations for PRIA FEP Fisheries..... 27

Table 4. Candidate ESA species, and ESA-listed species being evaluated for critical habitat designation 29

Table 5. Summary of habitat mapping in the PRIA..... 75

Table 6. Occurrence of EFH by feature in the PRIAs 77

Table 7. Mean percent cover of live coral from RAMP sites collected from towed-diver surveys in the PRIA 78

Table 8. Mean percent cover of macroalgae from RAMP sites collected from towed-diver surveys in the PRIA 78

Table 9. Mean percent cover of crustose coralline algae from RAMP sites collected from towed-diver surveys in the PRIA..... 79

Table 10. Level of EFH information available for the Western Pacific precious coral MUS 80

Table 11. Level of EFH information available for the Western Pacific BMUS and seamount groundfish MUS complex..... 80

Table 12. Level of EFH information available for the Western Pacific CMUS complex 81

Table 13. MMAs established under FEPs from 50 CFR § 665 87

Table 14. List of brainstormed potential archipelagic island fishery relationships scored and ranked from highest to lowest priority..... 90

TABLE OF FIGURES

Figure 1. Mean fish biomass ($\text{g/m}^2 \pm$ standard error) of CREMUS grouped by U.S. Pacific reef area from the years 2010 to 2018 by latitude; figure continued from previous page	5
Figure 2. Mean fish biomass ($\text{g/m}^2 \pm$ standard error) of PRIA CREMUS from 2010 to 2018 with American Samoa archipelago mean estimates overlaid (red lines); figure on from next page	7
Figure 3. Mean fish size (cm, TL \pm standard error) of PRIA CREMUS from 2010 to 2018 with American Samoa archipelago mean estimates overlaid (red lines); figure continued on next page	10
Figure 4. Settlement of the Pacific Islands, courtesy Wikimedia Commons. (from https://commons.wikimedia.org/wiki/File:Polynesian_Migration.svg).....	31
Figure 5. Indicators of change of pelagic coastal and marine systems; conceptual model	36
Figure 6. Schematic diagram illustrating how indicators are connected to one another and how they vary as a result of natural climate variability	37
Figure 7. Schematic diagram illustrating how indicators are connected to one another and how they vary as a result of anthropogenic climate change	38
Figure 8. Regional spatial grids representing the scale of the climate change indicators being monitored	38
Figure 9. Monthly mean (red) and seasonally-corrected (black) atmospheric carbon dioxide (ppm) at Mauna Loa Observatory, Hawai`i.....	40
Figure 10. Oceanic pH (black) and long-term trend (red) at Station ALOHA from 1989-2017..	41
Figure 11. Oceanic Niño Index from 1950-2018 (top) and 2000–2018 (bottom) with El Niño periods in red and La Niña periods in blue	42
Figure 12. Pacific Decadal Oscillation from 1950–2018 (top) and 2000–2018 (bottom) with positive warm periods in red and negative cool periods in blue.....	43
Figure 13. 2018 Pacific basin tropical cyclone tracks	46
Figure 14. Tropical storm totals by region.....	47
Figure 15. Accumulated Cyclone Energy (ACE) Index by region.....	47
Figure 16. Sea surface temperature climatology and anomalies from the PRIA Grid	49
Figure 17. Sea surface temperature climatology and anomalies from Johnston Atoll Grid.....	50
Figure 18. Sea surface temperature climatology and anomalies from Wake Atoll Grid.....	51
Figure 19. Coral Thermal Stress Exposure, Howland/Baker Virtual Station 2013-2018 (Coral Reef Watch Degree Heating Weeks)	53
Figure 20. Coral Thermal Stress Exposure, Northern Line Islands Virtual Station 2013-2018 (Coral Reef Watch Degree Heating Weeks).....	53
Figure 21. Coral Thermal Stress Exposure, Johnston Virtual Station 2013-2018 (Coral Reef Watch Degree Heating Weeks).....	54
Figure 22. Coral Thermal Stress Exposure, Wake Atoll Virtual Station 2013-2018 (Coral Reef Watch Degree Heating Weeks).....	54

Figure 23. Chlorophyll-A (Chl-A) and Chl-A Anomaly from the PRIA Grid 57

Figure 24. Chlorophyll-A (Chl-A) and Chl-A Anomaly from the Johnston Atoll Grid..... 58

Figure 25. Chlorophyll-A (Chl-A) and Chl-A Anomaly from the Wake Atoll Grid..... 59

Figure 26. CMAP precipitation across the PRIA Grid with 2018 values in red..... 61

Figure 27. CMAP precipitation across the Johnston Atoll Grid with 2018 values in red 61

Figure 28. CMAP precipitation across the Wake Atoll Grid with 2018 values in red 62

Figure 29a. Sea surface height and anomaly across the Pacific Ocean 63

Figure 30. Monthly mean sea level without the regular seasonal fluctuations due to coastal ocean temperatures, salinities, winds, atmospheric pressures, and ocean currents..... 65

Figure 31. Monthly mean sea level without the regular seasonal fluctuations due to coastal ocean temperatures, salinities, winds, atmospheric pressures, and ocean currents..... 66

Figure 32. Pacific Remote Island Areas and the associated Pacific Remote Islands Marine National Monument 73

Figure 33. The Substrate EFH Limit and 700 meter isobath around the PRIAs (from Ryan et al. 2009) 74

Figure 34. PRIA Land and Seafloor Primary Data Coverage (from Miller et al., 2011) 76

Figure 35. Regulated fishing areas of the PRIAs..... 86

ACRONYMS AND ABBREVIATIONS

Acronym	Meaning
ABC	Acceptable Biological Catch
ACL	Annual Catch Limits
AM	Accountability Measures
BiOp	Biological Opinion
BOEM	Bureau of Ocean Energy Management
BRFA	Bottomfish Restricted Fishing Area
BSIA	Best Scientific Information Available
CFR	Code of Federal Regulations
CMS	Coastal and Marine Spatial
CNMI	Commonwealth of the Northern Mariana Islands
Council	Western Pacific Regional Fishery Management Council
CPUE	Catch Per Unit Effort
CREMUS	Coral Reef Ecosystem Management Unit Species
CREP	Coral Reef Ecosystem Program (PIFSC)
EC	Ecosystem Component
ECS	Ecosystem Component Species
EEZ	Exclusive Economic Zone
EFH	Essential Fish Habitat
EO	Executive Order
ESA	Endangered Species Act
FEP	Fishery Ecosystem Plan
FMP	Fishery Management Plan
FR	Federal Register
GMRT	Global Multi-Resolution Topography Data Synthesis
HAPC	Habitat Area of Particular Concern
ITS	Incidental Take Statement
LOF	List of Fisheries
MBTA	Migratory Bird Treaty Act
MFMT	Maximum Fishing Mortality Threshold
MHI	Main Hawaiian Islands
MLCD	Marine Life Conservation District
MMA	Marine Managed Area
MMPA	Marine Mammal Protection Act
MPA	Marine Protected Area
MPCC	Marine Planning and Climate Change
MPCCC	Council's MPCC Committee
MSA	Magnuson-Stevens Fishery Conservation and Management Act
MSST	Minimum Stock Size Threshold
MSY	Maximum Sustainable Yield
MUS	Management Unit Species
MSFCMA	Magnuson-Stevens Fishery Conservation and Management Act

NEPA	National Environmental and Policy Act
nm	Nautical Miles
NMFS	National Marine Fisheries Service
NWHI	Northwestern Hawaiian Islands
ONI	Oceanic Niño Index
OFL	Overfishing Limit
OY	Optimum Yield
PDO	Pacific Decadal Oscillation
Pelagic FEP	Fishery Ecosystem Plan for the Pacific Pelagic Fisheries
PI	Pacific Islands
PIFSC	Pacific Islands Fisheries Science Center
PIRO	NOAA NMFS Pacific Islands Regional Office
PMUS	Pelagic Management Unit Species
PRIA	Pacific Remote Island Area
PRIMNM	Pacific Remote Islands Marine National Monument
RAMP	Reef Assessment and Monitoring Program (CREP)
ROA	Risk of Overfishing Analysis
RPB	Regional Planning Body
SAFE	Stock Assessment and Fishery Evaluation
SDC	Status Determination Criteria
SEEM	Social, Ecological, Economic, and Mgmt. (Uncertainty Analysis)
TAC	Total Annual Catch
USACE	United States Army Corps of Engineers
WPRFMC	Western Pacific Regional Fishery Management Council

This page was intentionally left blank.

1 FISHERY PERFORMANCE

Fisheries in the Pacific Remote Island Areas (PRIAs), including Palmyra Atoll, Kingman Reef, Jarvis Island, Baker Island, Howland Island, Johnston Atoll, and Wake Island, are limited. Fishery performance will be made available for the PRIAs in future reports as resources allow.

1.1 NUMBER OF FEDERAL PERMIT HOLDERS

The Code of Federal Regulations, Title 50 Part 665 requires the following Federal permits for fishing in the exclusive economic zone (EEZ) of the PRIAs.

1.1.1 Special Coral Reef Ecosystem Permit

Regulations require a special coral reef ecosystem fishing permit for anyone fishing for coral reef management unit species (CREMUS) in a low-use marine protected area (MPA), fishing for species on the list of Potentially Harvested Coral Reef Taxa, or using fishing gear not specifically allowed in the regulations. The National Marine Fisheries Service (NMFS) will make an exception to this permit requirement for any person issued a permit to fish under any fishery ecosystem plan (FEP) who incidentally catches a CREMUS while fishing for bottomfish management unit species (BMUS), crustacean management unit species (CMUS), western Pacific pelagic management unit species (PMUS), precious coral (PCMUS), or seamount groundfish. Regulations require a transshipment permit for any receiving vessel used to land or transship potentially harvested coral reef taxa, or any CREMUS caught in a low-use MPA.

1.1.2 Western Pacific Precious Corals Permit

Regulations require a Western Pacific Precious Corals permit for anyone harvesting or landing black, bamboo, pink, red, or gold corals in the EEZs of the U.S. Western Pacific.

1.1.3 Western Pacific Crustaceans Permit (Lobster or Deepwater Shrimp)

Regulations require a Western Pacific Crustaceans permit for any owner of a U.S. fishing vessel used to fish for lobster or deepwater shrimp in the EEZs around of the U.S. Western Pacific.

1.1.4 PRIA Bottomfish Permit

Regulations require obtaining a PRIA Bottomfish permit for anyone using bottomfish gear to fish for BMUS in the EEZ around the PRIAs. Commercial fishing is prohibited within the boundaries of the Pacific Remote Islands Marine National Monument (PRIMNM).

There is no record of coral reef or precious coral fishery permits issued for the EEZ around the PRIAs since 2008. Table 1 provides the number of permits issued for PRIA fisheries from 2009 to 2018. Historical data from were accessed from PIFSC, and data for 2018 are from the PIRO Sustainable Fisheries Division (SFD) permits program.

Table 1. Number of federal permit holders in the lobster, shrimp, and bottomfish fisheries of the PRIA from 2009 to 2018

PRIA Fisheries	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
Lobster	3	0	0	0	0	0	0	0	0	0
Shrimp	0	1	0	0	0	0	0	0	0	0
Bottomfish	3	6	5	4	1	2	0	1	1	4

1.2 ADMINISTRATIVE AND REGULATORY ACTIONS

This summary describes management actions for PRIA fisheries that NMFS implemented after the April 2018 Joint FEP Plan Team meeting.

June 14, 2018. Final rule. 5-Year Extension of Moratorium on Harvest of Gold Corals. This final rule extends the region-wide moratorium on the harvest of gold corals in the U.S. Pacific Islands through June 30, 2023. NOAA Fisheries intends this final rule to prevent overfishing and to stimulate research on gold corals.

February 8, 2019. Final rule. Reclassifying Management Unit Species to Ecosystem Component Species. This final rule reclassifies certain management unit species in the Pacific Islands as ecosystem component species. The rule also updates the scientific and local names of certain species. The intent of this final rule is to prioritize conservation and management efforts and to improve efficiency of fishery management in the region. This rule is effective March 11, 2019.

2 ECOSYSTEM CONSIDERATIONS

2.1 CORAL REEF FISH ECOSYSTEM PARAMETERS

2.1.1 Regional Reef Fish Biomass

Description: ‘Reef fish biomass’ is mean biomass of coral reef fishes per unit area derived from visual survey data between 2010 and 2018. These data are shown in **Error! Reference source not found.**

Rationale: Reef fish biomass has been widely used as an indicator of relative ecosystem status, and has repeatedly been shown to be sensitive to changes in fishing pressure, habitat quality, and oceanographic regime.

Data Category: Fishery-independent


Timeframe: Triennial

Jurisdiction: American Samoa, Guam, CNMI, MHI, NWHI, and the PRIAs

Spatial Scale: Regional

Data Source: Data used to generate biomass estimates comes from visual surveys conducted by NOAA PIFSC Coral Reef Ecosystem and partners, as part of the Pacific Reef Assessment and Monitoring Program (RAMP) (http://www.pifsc.noaa.gov/cred/pacific_ramp.php). Survey methods are described in detail in Ayotte et al. (2015), but in brief involve teams of divers conducting stationary point count cylinder (SPC) surveys within a target domain of <30 meter hard-bottom habitat at each island, stratified by depth zone and, for larger islands, by section of coastline. For consistency among islands, only data from forereef habitats are used. At each SPC, divers record the number, size, and species of all fishes within or passing through paired 15 meter-diameter cylinders over the course of a standard count procedure.

Fish sizes and abundance are converted to biomass using standard length-to-weight conversion parameters, taken largely from FishBase (<http://www.fishbase.org>), and converted to biomass per unit area by dividing by the area sampled per survey. Site-level data were pooled into island-scale values by first calculating mean and variance within strata, and then calculating weighted island-scale mean and variance using the formulas given in Smith et al. (2011), with strata weighted by their respective sizes.






Figure 1. Mean fish biomass (g/m² ± standard error) of CREMUS grouped by U.S. Pacific reef area from the years 2010 to 2018 by latitude; figure continued from previous page

2.1.2 Archipelagic Reef Fish Biomass

Description: ‘Reef fish biomass’ is mean biomass of coral reef fishes per unit area derived from visual survey data between 2010 and 2018. These data are shown in Figure 2.

Rationale: Identical to the rationale described in Section 2.1.1.

Data Category: Fishery-independent

Timeframe: Triennial

Jurisdiction: PRIAs

Spatial Scale: Island

Data Source: Data used to generate biomass estimates comes from visual surveys conducted by NOAA PIFSC Coral Reef Ecosystem Division (CRED) and partners, as part of the Pacific RAMP. Survey methods and sampling design, and methods to generate reef fish biomass are described above (Section 2.1.1).





Figure 2. Mean fish biomass (g/m² ± standard error) of PRIA CREMUS from 2010 to 2018 with American Samoa archipelago mean estimates overlaid (red lines); figure continued on next page



2.1.3 Archipelagic Mean Fish Size

Description: ‘Mean fish size’ is mean size of reef fishes > 10 cm total length (TL), excluding small fishes, derived from visual survey data between 2010 and 2018. These data are shown in Figure 3.

Rationale: Mean size is important as it is widely used as an indicator of fishing pressure. A fishery can sometimes preferentially target large individuals, and can also the number of fishes reaching older (and larger) size classes. Large fishes contribute disproportionately to community fecundity and can have important ecological roles; for example, excavating bites by large parrotfishes probably have a longer lasting impact on reef benthos than bites by smaller fishes.

Data Category: Fishery-independent

Timeframe: Triennial

Jurisdiction: PRIAs

Spatial Scale: Island

Data Source: Data used to generate biomass estimates comes from visual surveys conducted by NOAA PIFSC Coral Reef Ecosystem and partners, as part of the Pacific RAMP. Survey methods and sampling design, and methods to generate reef fish biomass are described above (Section 2.1.1). Fishes smaller than 10 cm TL are excluded so that the fish assemblage measured more closely reflects fishes that are potentially fished, and so that mean sizes are not overly influenced by variability in space and time of recent recruitment.





Figure 3. Mean fish size (cm, TL \pm standard error) of PRIA CREMUS from 2010 to 2018 with American Samoa archipelago mean estimates overlaid (red lines); figure continued on next page



2.1.4 Reef Fish Population Estimates

Description: ‘Reef fish population estimates’ are calculated by multiplying mean biomass per unit area by estimated hard bottom area in a consistent habitat across all islands (specifically, the area of hard bottom forereef habitat in < 30 meters of water). These data are shown in Table 2.

Rationale: Reef fish population estimate data have utility in understanding the size of populations from which fishery harvests are extracted.

Data Category: Fishery-independent

Timeframe: Triennial

Jurisdiction: PRIAs

Spatial Scale: Island

Data Source: Data used to generate mean size estimates come from visual surveys conducted by NOAA PIFSC Coral Reef Ecosystem and partners, as part of the Pacific RAMP. Survey methods and sampling design, and methods to generate reef fish biomass are described above (Section 2.1.1). Those estimates are converted to population estimates by multiplying biomass (g/m^2) per island by the estimated area of hard bottom habitat < 30 meters deep at the island, which is the survey domain for the monitoring program that biomass data comes from. Measures of estimated habitat area per island are derived from Geographic Information System (GIS) bathymetry and NOAA Coral Reef Ecosystems Program (CREP) habitat maps.

Many reef fish taxa are present in other habitats than is surveyed by the program, and some taxa likely have the majority of their populations in deeper water. Additionally, fish counts have the potential to be biased by the nature of fish response to divers. Curious fishes, particularly in locations where divers are not perceived as a threat, will tend to be overestimated by visual survey, while skittish fishes will tend to be undercounted. It is also likely that numbers of jacks and sharks in some locations, such as the NWHI are overestimated by visual survey. Nevertheless, the data shown here are consistently gathered across space and time.

Table 2. Reef fish population estimates for PRIA CREMUS in 0-30 m hard bottom habitat

Island/Atoll	Total area of reef (Ha)	N	Estimated population biomass (metric tons) in survey domain of < 30 m hard bottom					
			Acanthuridae	Carangidae	Carcharhinids	Holocentridae	Kyphosidae	Labridae
Wake	279.8	128	13.2	19.3	1.7	4.9	31.5	7.0
Johnston	9,805.5	104	1,302.9	735.3	81.9	199.4	17.2	171.4
Kingman	3,751.3	171	365.8	113.1	1,924.3	59.7	-	89.6
Palmyra	2,812.0	210	318.6	233.4	716.5	32.6	3.3	54.1
Howland	172.9	119	22.3	14.0	26.9	12.5	0.8	2.2
Baker	390.3	113	54.8	42.3	71.1	24.7	2.4	6.9
Jarvis	365.9	231	78.3	45.7	162.3	17.5	3.0	15.1
TOTAL	17,577.6	1,076	2,002.8	1,155.3	3,709.5	319.2	72.8	344.4
Island/Atoll	Total area of reef (Ha)	N	Lethrinidae	Lutjanidae	Mullidae	Scaridae	Serranidae	<i>C. undulatus</i>
Wake	279.8	128	2.2	2.6	3.2	23.8	9.8	6.9
Johnston	9,805.5	104	35.8	375.2	119.9	673.0	-	-
Kingman	3,751.3	171	72.0	1,022.8	18.9	487.0	167.4	-
Palmyra	2,812.0	210	103.1	614.4	18.1	272.2	157.6	135.6
Howland	172.9	119	0.3	19.1	2.3	6.6	14.0	-
Baker	390.3	113	1.4	42.3	3.3	25.3	21.8	-
Jarvis	365.9	231	3.3	69.9	5.9	48.3	27.3	1.1
TOTAL	17,577.6	1,076	276.2	2,656.1	148.0	1,618.5	550.3	205.6

Notes: (1) N is the number of sites surveyed.

(2) No Siganidae or *Bolbometopon muricatum* were observed in the PRIAs during these surveys.

2.1.5 References

- Ayotte, P., McCoy, K., Heenan, A., Williams, I., and J. Zamzow. 2015. Coral Reef Ecosystem Division standard operating procedures: data collection for Rapid Ecological Assessment fish surveys. Retrieved from http://www.pifsc.noaa.gov/library/pubs/admin/PIFSC_Admin_Rep_15-07.pdf.
- Smith, S.G., Ault, J.S., Bohnsack, J.A., Harper, D.E., Luo, J.G., and D.B. McClellan. 2011. Multispecies survey design for assessing reef-fish stocks, spatially explicit management performance, and ecosystem condition. *Fisheries Research*, 109(1), p. 25-41. <http://doi.org/10.1016/j.fishres.2011.01.012>.

2.2 PROTECTED SPECIES

This section of the report summarizes information on protected species interactions in fisheries managed under the PRIA FEP. Protected species covered in this report include sea turtles, seabirds, marine mammals, elasmobranchs, and precious corals. Most of these species are protected under the Endangered Species Act (ESA), the Marine Mammal Protection Act (MMPA), and/or the Migratory Bird Treaty Act (MBTA). A list of protected species found in or near PRIA waters and a list of critical habitat designations in the Pacific Ocean are included in Appendix B.

2.2.1 Monitoring Protected Species Interactions in the PRIA FEP Fisheries

This report monitors the status of protected species interactions in the PRIA FEP fisheries using proxy indicators such as fishing effort and changes in gear types as these fisheries do not have observer coverage. Logbook programs are not expected to provide reliable data about protected species interactions due to the lack of active fisheries in these areas.

2.2.1.1 FEP Conservation Measures

Bottomfish, precious coral, coral reef, and crustacean fisheries managed under this FEP have not had reported interactions with protected species, and no specific regulations are in place to mitigate protected species interactions. Destructive gear such as bottom trawls, bottom gillnets, explosives, and poisons are prohibited under this FEP, and these prohibitions benefit protected species by preventing potential interactions with non-selective fishing gear.

2.2.1.2 ESA Consultations

ESA consultations were conducted by NMFS and the U.S. Fish and Wildlife Service (USFWS; for species under their jurisdiction) to ensure ongoing fisheries operations managed under the PRIA FEP are not jeopardizing the continued existence of any listed species or adversely modifying critical habitat. The results of these consultations, conducted under section 7 of the ESA, are briefly described below and summarized in Table 3.

NMFS concluded in an informal consultation dated February 20, 2015 that all fisheries managed under the PRIA FEP are not likely to adversely affect the Indo-West Pacific DPS of scalloped hammerhead shark. NMFS concluded on January 16, 2015 that all fisheries managed under the PRIA FEP have no effects on ESA-listed reef-building corals.

In January 2018, oceanic whitetip sharks and giant manta rays were listed under the ESA (83 FR 4153 and 83 FR 2916, respectively). If NMFS determines that the PRIA fisheries are likely to adversely affect these species, NMFS will initiate consultation for these two species for the applicable fisheries.

Table 3. Summary of ESA consultations for PRIA FEP Fisheries

Fishery	Consultation Date	Consultation Type^a	Outcome^b	Species
Bottomfish	3/8/2002	BiOp	NLAA	Loggerhead sea turtle, leatherback sea turtle, olive ridley sea turtle, green sea turtle, hawksbill sea turtle, humpback whale, blue whale, fin whale, sei whale, sperm whale
Coral reef ecosystem	3/7/2002	LOC	NLAA	Loggerhead sea turtle, leatherback sea turtle, olive ridley sea turtle, green sea turtle, hawksbill sea turtle, humpback whale, blue whale, fin whale, sei whale, sperm whale
	5/22/2002	LOC (USFWS)	NLAA	Green, hawksbill, leatherback, loggerhead and olive ridley turtles, Newell's shearwater, short-tailed albatross, Laysan duck, Laysan finch, Nihoa finch, Nihoa millerbird, Micronesian megapode, 6 terrestrial plants
Crustacean	9/28/2007	LOC	NLAA	Loggerhead sea turtle, leatherback sea turtle, olive ridley sea turtle, green sea turtle, hawksbill sea turtle, humpback whale, blue whale, fin whale, sei whale, sperm whale
Precious coral	10/4/1978	BiOp	Does not constitute threat	Sperm whale, leatherback sea turtle
	12/20/2000	LOC	NLAA	Humpback whale, green sea turtle, hawksbill sea turtle
All fisheries	1/16/2015	No effect memo	No effect	Reef-building corals
	2/20/2015	LOC	NLAA	Scalloped hammerhead shark (Indo-west Pacific DPS)

^a BiOp = Biological Opinion; LOC = Letter of Concurrence

^b LAA = likely to adversely affect; NLAA = not likely to adversely affect.

Bottomfish Fishery

In a biological opinion issued on March 3, 2002, NMFS concluded that the ongoing operation of the Western Pacific Region's bottomfish and seamount fisheries is not likely to jeopardize the continued existence of five sea turtle species (loggerhead, leatherback, olive ridley, green, and hawksbill turtles) and five marine mammal species (humpback, blue, fin, sei, and sperm whales)

Crustacean Fishery

An informal consultation completed by NMFS on September 28, 2007 concluded that PRIA crustacean fisheries are not likely to adversely affect five sea turtle species (loggerhead, leatherback, olive ridley, green, and hawksbill turtles) and five marine mammal species (humpback, blue, fin, sei, and sperm whales).

Coral Reef Fishery

An informal consultation completed by NMFS on March 7, 2002 concluded that fishing activities conducted under the Coral Reef Ecosystems FMP are not likely to adversely affect five

sea turtle species (loggerhead, leatherback, olive ridley, green, and hawksbill turtles) and five marine mammal species (humpback, blue, fin, sei, and sperm whales).

On May 22, 2002, the USFWS concurred with the determination of NMFS that the activities conducted under the Coral Reef Ecosystems FMP are not likely to adversely affect listed species under USFWS's exclusive jurisdiction (i.e., seabirds and terrestrial plants) and listed species shared with NMFS (i.e., sea turtles).

Precious Coral Fishery

An informal consultation completed by NMFS on December 20, 2000 concluded that PRIA precious coral fisheries are not likely to adversely affect humpback whales, green turtles, or hawksbill turtles.

2.2.1.3 Non-ESA Marine Mammals

The MMPA requires NMFS to annually publish a List of Fisheries (LOF) that classifies commercial fisheries in one of three categories based on the level of mortality and serious injury of marine mammals associated with that fishery. PRIA fisheries are not classified under the LOF due to the lack of active commercial fisheries.

2.2.2 Status of Protected Species Interactions in the PRIA FEP Fisheries There are currently no bottomfish, crustacean, coral reef, or precious coral fisheries operating in the PRIA and no historical observer data are available for fisheries under this FEP. No new fishing activity has been reported, and there is no other information to indicate that impacts to protected species from PRIA fisheries have changed in recent years.

2.2.3 Identification of Emerging Issues

Several ESA-listed species are being evaluated for critical habitat designation (Table 4). If critical habitats are designated, they will be included in this SAFE report and impacts from FEP-managed fisheries will be evaluated under applicable mandates.

Table 4. Candidate ESA species, and ESA-listed species being evaluated for critical habitat designation

Species		Listing Process			Post-Listing Activity	
Common Name	Scientific Name	90-Day Finding	12-Month Finding / Proposed Rule	Final Rule	Critical Habitat	Recovery Plan
Oceanic whitetip shark	<i>Carcharhinus longimanus</i>	Positive (81 FR 1376, 1/12/2016)	Positive, threatened (81 FR 96304, 12/29/2016)	Listed as Threatened (83 FR 4153, 1/30/18)	Not determinable because of insufficient data (83 FR 4153, 1/30/18)	TBA
Pacific bluefin tuna	<i>Thunnus orientalis</i>	Positive (81 FR 70074, 10/11/2016)	Not warranted (82 FR 37060, 8/8/17)	N/A	N/A	N/A
Giant manta ray	<i>Manta birostris</i>	Positive (81 FR 8874, 2/23/2016)	Positive, threatened (82 FRN 3694, 1/12/2017)	Listed as Threatened (83 FR 2916, 1/22/18)	N/A	N/A
Corals	N/A	Positive for 82 species (75 FR 6616, 2/10/2010)	Positive for 66 species (77 FR 73219, 12/7/2012)	20 species listed as threatened (79 FR 53851, 9/10/2014)	In development, proposal expected TBA	In development, expected TBA, interim recovery outline in place
Cauliflower coral	<i>Pocillopora meandrina</i>	Positive (83 FR 47592, 9/20/2018)	TBD	TBD	N/A	N/A
Giant Clams	<i>Hippopus hippopus</i> , <i>H. porcellanus</i> , <i>Tridacna costata</i> , <i>T. derasa</i> , <i>T. gigas</i> , <i>T. squamosa</i> , and <i>T. tevoroa</i>	Positive (82 FR 28946, 06/26/2017)	TBD (status review ongoing)	TBD	N/A	N/A
Green sea turtle	<i>Chelonia mydas</i>	Positive (77 FR 45571, 8/1/2012)	Identification of 11 DPSs, endangered and threatened (80 FR 15271, 3/23/2015)	11 DPSs listed as endangered and threatened (81 FR 20057, 4/6/2016)	In development, proposal expected TBA ^a	TBA

Leatherback sea turtle	<i>Dermochelys coriacea</i>	Positive 90-day finding on a petition to identify the Northwest Atlantic leatherback turtle as a DPS (82 FR 57565, 12/06/2017)	TBA (status review ongoing)	TBA	N/A	N/A
------------------------	-----------------------------	--	-----------------------------	-----	-----	-----

^a NMFS and USFWS have been tasked with higher priorities regarding sea turtle listings under the ESA, and do not anticipate proposing green turtle critical habitat designations in the immediate future.

2.2.4 Identification of Research, Data, and Assessment Needs

The following research, data, and assessment needs for insular fisheries were identified by the Council’s Protected Species Advisory Committee and Plan Team:

- Improve the precision of commercial and non-commercial fisheries data to improve understanding of potential protected species impacts.
- Define and evaluate innovative approaches to derive robust estimates of protected species interactions in insular fisheries.

2.3 SOCIOECONOMICS

This section outlines the pertinent economic, social, and community information available for assessing the successes and impacts of management measures and the achievements of the Fishery Ecosystem Plan (FEP) for the Pacific Remote Island Area (PRIA; WPRFMC, 2018). It meets the objective of “Support Fishing Communities” adopted at the 165th Council meeting; specifically, it identifies the various social and economic groups within the regions’ fishing communities and their interconnections. The section begins with an overview of the socioeconomic context for the region, and then provides a summary of relevant studies and data for the PRIA.

In 1996, the Magnuson-Stevens Fishery Conservation and Management Act’s National Standard 8 (NS8) specified that conservation and management measures need to account for the importance of fishery resources in fishing communities, to support sustained participation in the fisheries, and to minimize adverse economic impacts, provided that these considerations do not compromise conservation. Unlike other regions of the U.S., the settlement of the Western Pacific region was intimately tied to the ocean, which is reflected in local culture, customs, and traditions (Figure 4).



Figure 4. Settlement of the Pacific Islands, courtesy Wikimedia Commons. (from https://commons.wikimedia.org/wiki/File:Polynesian_Migration.svg)

Polynesian voyagers relied on the ocean and marine resources on their long voyages in search of new islands, as well as in sustaining established island communities. Today, the population of the region also represents many Asian cultures from Pacific Rim countries, which have a similar reliance on marine resources. Thus, fishing and seafood are integral to local community ways of life. This is reflected in the amount of seafood eaten in the region relative to the rest of the United States, as well as in the language, customs, ceremonies, and community events. The amount of available seafood can also affect seasonality in prices of fish. Because fishing is such an integral part of the culture, it is difficult to discern commercial from non-commercial fishing where most trips involving multiple motivations and multiple uses of the fish caught. While the economic perspective is an important consideration, fishermen report other motivations, such as customary exchange, as being equally important. Due to changing economies and westernization, waning recruitment of younger fishermen is becoming a concern for the sustainability of fishing and fishing traditions in the region.

2.3.1 Response to Previous Council Recommendations

There were no Council Recommendations related to socioeconomic considerations in the PRIA during 2018.

2.3.2 Background

Human habitation in the PRIA is limited. The FEP for the PRIA provides a description of the geography, history, and socioeconomic considerations of the archipelago (WPRFMC 2018). Grace-McCaskey (2014) provided a brief review of the importance of these areas from a cultural perspective. She noted that although the PRIA were uninhabited when first visited by Westerners, Polynesians and Micronesians likely had been periodically visiting these islands for centuries. Many of the islands in the PRIA were altered during WWII, and many have subsequently become National Wildlife Refuges or part of the Pacific Remote Islands Marine National Monument. Only Wake, Johnston, and Palmyra have seasonal- and year-round residents, primarily related to the U.S. military and refuge management. The surrounding reef ecosystems are considered to be some of the healthiest in the world due to their distance to areas of high human population densities, though some are experiencing residual impacts from military activity nearby. There are no designated fishing communities residing in the PRIA. Most of the fishing effort has been concentrated around Johnston and Palmyra by members of the Hawaii fishing community.

2.3.3 Ongoing Research and Information Collection

There is currently no ongoing research specific to the PRIA.

2.3.4 Relevant PIFSC Economics and Human Dimensions Publications: 2018

There were no relevant PIFSC publications regarding the economics or human dimensions of the PRIAs in 2018.

2.3.5 References

- Grace-McCaskey, C. 2014. Examining the potential of using secondary data to better understand human-reef relationships across the Pacific. Pacific Islands Fish. Sci. Cent., Natl. Mar. Fish. Serv., NOAA, Honolulu, HI 96818-5007. Pacific Islands Fish. Sci. Cent. Admin. Rep. H-14-01, 69 p.
https://www.pifsc.noaa.gov/library/pubs/admin/PIFSC_Admin_Rep_14-01.pdf
- Polovina, J. and Dreftak, K., (Chairs), Baker, J., Bloom, S., Brooke, S., Chan, V., Ellgen, S., Golden, D., Hospital, J., Van Houtan, K., Kolinski, S., Lumsden, B., Maison, K., Mansker, M., Oliver, T., Spalding, S., and P. Woodworth-Jefcoats, 2016. Pacific Islands Regional Action Plan: NOAA Fisheries climate science strategy. U.S. Dept. of Commerce, NOAA Technical Memorandum NOAA-TM-NMFS-PIFSC-59, 33 p. doi:10.7289/V5/TM-PIFSC-59.
- WPRFMC, 2018. Annual Stock Assessment and Fishery Evaluation Report: Pacific Remote Island Area Fishery Ecosystem Plan 2017. Sabater, M., Ishizaki, A., Remington, T., Spalding, S. (Eds.) Western Pacific Regional Fishery Management Council. Honolulu, Hawaii 96813 USA.

2.4 CLIMATE AND OCEANIC INDICATORS

2.4.1 Introduction

Over the past few years, the Council has incorporated climate change into the overall management of the fisheries over which it has jurisdiction. This 2018 Annual SAFE Report includes a now standard chapter on indicators of climate and oceanic conditions in the Western Pacific region. These indicators reflect global climate variability and change as well as trends in local oceanographic conditions.

The reasons for the Council's decision to provide and maintain an evolving discussion of climate conditions as an integral and continuous consideration in their deliberations, decisions, and reports are numerous:

- Emerging scientific and community understanding of the impacts of changing climate conditions on fishery resources, the ecosystems that sustain those resources, and the communities that depend upon them;
- Recent Federal Directives including the 2010 implementation of a National Ocean Policy that identified Resiliency and Adaptation to Climate Change and Ocean Acidification as one of nine National priorities as well as the development of a Climate Science Strategy by NMFS in 2015 and the subsequent development of the Pacific Islands Regional Action Plan for climate science; and
- The Council's own engagement with NOAA as well as jurisdictional fishery management agencies in American Samoa, CNMI, Guam, and Hawai'i as well as fishing industry representatives and local communities in those jurisdictions.

In 2013, the Council began restructuring its Marine Protected Area/Coastal and Marine Spatial Planning Committee to include a focus on climate change, and the committee was renamed as the Marine Planning and Climate Change (MPCC) Committee. In 2015, based on recommendations from the committee, the Council adopted its Marine Planning and Climate Change Policy and Action Plan, which provided guidance to the Council on implementing climate change measures, including climate change research and data needs. The revised Pelagic Fisheries Ecosystem Plan (FEP; February 2016) included a discussion on climate change data and research as well as a new objective (Objective 9) that states the Council should consider the implications of climate change in decision-making, with the following sub-objectives:

- a) To identify and prioritize research that examines the effects of climate change on Council-managed fisheries and fishing communities.
- b) To ensure climate change considerations are incorporated into the analysis of management alternatives.
- c) To monitor climate change related variables via the Council's Annual Reports.
- d) To engage in climate change outreach with U.S. Pacific Islands communities.

Beginning with the 2015 report, the Council and its partners began providing continuing descriptions of changes in a series of climate and oceanic indicators.

This annual report focuses previous years' efforts by refining existing indicators and improving communication of their relevance and status. Future reports will include additional indicators as the information becomes available and their relevance to the development, evaluation, and revision of the FEPs becomes clearer. Working with national and jurisdictional partners, the Council will make all datasets used in the preparation of this and future reports available and easily accessible. **Response to Previous Council Recommendations**

At its 170th meeting from June 20-22, 2017, the Council directed staff to support the development of community training and outreach materials and activities on climate change. In addition, the Council directed staff to coordinate a “train-the-trainers” workshop that includes NOAA scientists who presented at the 6th Marine Planning and Climate Change Committee (MPCCC) meeting and the MPCCC committee members in preparation for community workshops on climate and fisheries. The Council and NOAA partnered to deliver the workshops in the fall of 2017 to the MPCCC members in Hawaii (with the Hawaii Regional Ecosystem Advisory Committee), as well as American Samoa, Guam, and the CNMI (with their respective Advisory Panel groups). Feedback from workshop participants has been incorporated into this year's climate and oceanic indicator section. To prepare for community outreach, Guam-based MPCCC members conducted a climate change survey and shared the results with the MPCCC at its 7th meeting on April 10th and 11th, 2018. The Council also directed staff to explore funding avenues to support the development of additional oceanic and climate indicators, such as wind and extratropical storms. These indicators were added to this module by corresponding Plan Team members in 2018. There were no Council recommendations relevant to the climate and oceanic indicators section of the Annual SAFE Report in 2018.


Prior to holding its 8th meeting, the MPCCC was disbanded in early 2019, re-allocating its responsibilities among its members already on other committees or teams, such as the Fishery Ecosystem Plan Teams.

2.4.2 Conceptual Model

In developing this chapter, the Council relied on a number of recent reports conducted in the context of the U.S. National Climate Assessment including, most notably, the 2012 Pacific Islands Regional Climate Assessment and the Ocean and Coasts chapter of the 2014 report on a Pilot Indicator System prepared by the National Climate Assessment and Development Advisory Committee (NCADAC).

The Advisory Committee Report presented a possible conceptual framework designed to illustrate how climate factors can connect to and interact with other ecosystem components to impact ocean and coastal ecosystems and human communities. The Council adapted this model with considerations relevant to the fishery resources of the Western Pacific Region:

Indicators of Change to Pelagic Coastal and Marine Systems*
(Items in red to be monitored for 2015 Annual Report of the Pelagic Fishery Ecosystem Plan for the Western Pacific Region)



**Adapted from National Climate Assessment and Development Advisory Committee, February 2014. National Climate Indicators System Report, B-59.*

Figure 5. Indicators of change of pelagic coastal and marine systems; conceptual model

As described in the 2014 NCADAC report, the conceptual model presents a “simplified representation of climate and non-climate stressors in coastal and marine ecosystems.” For the purposes of this Annual Report, the modified Conceptual Model allows the Council and its partners to identify indicators of interest to be monitored on a continuing basis in coming years. The indicators shown in red were considered for inclusion in the Annual SAFE Reports, though the final list of indicators varied somewhat. Other indicators will be added over time as data become available and an understanding of the causal chain from stressors to impacts emerges.

The Council also hopes that this Conceptual Model can provide a guide for future monitoring and research. This guide will ideally enable the Council and its partners to move forward from observations and correlations to understanding the specific nature of interactions, and to develop capabilities to predict future changes of importance in the developing, evaluating, and adapting of FEPs in the Western Pacific region.

2.4.3 Selected Indicators

The primary goal for selecting the indicators used in this report is to provide fisheries-related communities, resource managers, and businesses with a climate-related situational awareness. In this context, indicators were selected to:

- Be fisheries relevant and informative.
- Build intuition about current conditions in light of a changing climate;
- Provide historical context; and
- Allow for recognition of patterns and trends.

In this context, this section includes the following climate and oceanic indicators:

- Atmospheric concentration of carbon dioxide (CO₂)
- Oceanic pH at Station ALOHA;
- Oceanic Niño Index (ONI);
- Pacific Decadal Oscillation (PDO);
- Tropical cyclones;
- Sea surface temperature (SST);
- Coral Thermal Stress Exposure;
- Chlorophyll-A;
- Rainfall; and
- Sea Level (Sea Surface Height).

Figures 2 and 3 provide a description of these indicators and illustrate how they are connected to each other in terms of natural climate variability and anthropogenic climate change.




Figure 6. Schematic diagram illustrating how indicators are connected to one another and how they vary as a result of natural climate variability




Figure 7. Schematic diagram illustrating how indicators are connected to one another and how they vary as a result of anthropogenic climate change




Figure 8. Regional spatial grids representing the scale of the climate change indicators being monitored

1.1.1.1 Atmospheric Concentration of Carbon Dioxide (CO₂) at Mauna Loa

Rationale: Atmospheric carbon dioxide is a measure of what human activity has already done to affect the climate system through greenhouse gas emissions. It provides quantitative information in a simplified, standardized format that decision makers can easily understand. This indicator demonstrates that the concentration (and, in turn, warming influence) of greenhouse gases in the atmosphere has increased substantially over the last several decades.

Status: Atmospheric CO₂ is increasing exponentially. This means that atmospheric CO₂ is increasing at a faster rate each year. In 2018, the annual mean concentration of CO₂ was 409 ppm. In 1959, the first year of the time series, it was 316 ppm. The annual mean passed 350 ppm in 1988 and 400 ppm in 2015.

Description: Monthly mean atmospheric carbon dioxide (CO₂) at Mauna Loa Observatory, Hawai`i in parts per million (ppm) from March 1958 to present. The observed increase in monthly average carbon dioxide concentration is primarily due to CO₂ emissions from fossil fuel burning. Carbon dioxide remains in the atmosphere for a very long time, and emissions from any location mix throughout the atmosphere in approximately one year. The annual variations at Mauna Loa, Hawai`i are due to the seasonal imbalance between the photosynthesis and respiration of terrestrial plants. During the summer growing season, photosynthesis exceeds respiration, and CO₂ is removed from the atmosphere. In the winter (outside the growing season), respiration exceeds photosynthesis, and CO₂ is returned to the atmosphere. The seasonal cycle is strongest in the northern hemisphere because of its larger land mass.

Timeframe: Annual, monthly.

Region/Location: Mauna Loa, Hawai`i, but representative of global atmospheric carbon dioxide concentration.

Measurement Platform: *In-situ* station.

Sourced from: NOAA ESRL (2019a).




Figure 9. Monthly mean (red) and seasonally-corrected (black) atmospheric carbon dioxide (ppm) at Mauna Loa Observatory, Hawai'i

1.1.1.2 Oceanic pH

Rationale: Oceanic pH is a measure of how greenhouse gas emissions have already impacted the ocean. This indicator demonstrates that oceanic pH has decreased significantly over the past several decades (i.e. the ocean has become more acidic). Increasing ocean acidification limits the ability of marine organisms to build shells and other calcareous structures. Recent research has shown that pelagic organisms such as pteropods and other prey for commercially-valuable fish species are already being negatively impacted by increasing acidification (Feely *et al.*, 2016). The full impact of ocean acidification on the pelagic food web is an area of active research (Fabry *et al.*, 2008).

Status: The ocean is roughly 9.4% more acidic than it was nearly 30 years ago at the start of this time series. Over this time, pH has declined by 0.0389 at a constant rate. In 2017, the most recent year for which data are available, the average pH was 8.07. Additionally, small variations seen over the course of the year are now outside the range seen in the first year of the time series. The highest pH value reported for the most recent year (8.0831) is lower than the lowest pH value reported in the first year of the time series (8.0845).

Description: Trends in surface (5 m) pH at Station ALOHA, north of Oahu (22.75°N, 158°W), collected by the Hawai'i Ocean Time Series (HOT) from October 1988 to 2017 (2018 data are not yet available). Oceanic pH is a measure of ocean acidity, which increases as the ocean absorbs carbon dioxide from the atmosphere. Lower pH values represent greater acidity. Oceanic pH is calculated from total alkalinity (TA) and dissolved inorganic carbon (DIC). Total alkalinity represents the ocean's capacity to resist acidification as it absorbs CO₂ and the amount of CO₂

absorbed is captured through measurements of DIC. The multi-decadal time series at Station ALOHA represents the best available documentation of the significant downward trend in oceanic pH since the time series began in 1988. Oceanic pH varies over both time and space, though the conditions at Station ALOHA are considered broadly representative of those across the Western and Central Pacific’s pelagic fishing grounds.

Timeframe: Monthly.

Region/Location: Station ALOHA: 22.75°N, 158°W.

Measurement Platform: *In-situ* station.

Sourced from: the Hawaii Ocean Time Series as described in Karl et al. (1996) and on its website (HOT, 2019).




Figure 10. Oceanic pH (black) and long-term trend (red) at Station ALOHA from 1989-2017

1.1.1.3 Oceanic Niño Index (ONI)

Rationale: The El Niño – Southern Oscillation (ENSO) cycle is known to have impacts on Pacific fisheries including tuna fisheries. The ONI focuses on ocean temperature, which has the most direct effect on these fisheries.

Status: In 2018, the ONI transitioned from a weak La Niña to neutral conditions.

Description: The three-month running mean of satellite remotely-sensed sea surface temperature (SST) anomalies in the Niño 3.4 region (5°S – 5°N, 120° – 170°W). The Oceanic Niño Index (ONI) is a measure of the El Niño – Southern Oscillation (ENSO) phase. Warm and cool phases, termed El Niño and La Niña respectively, are based in part on an ONI threshold of ± 0.5 °C being met for a minimum of five consecutive overlapping seasons. Additional atmospheric indices are needed to confirm an El Niño or La Niña event, as the ENSO is a coupled ocean-atmosphere phenomenon. The atmospheric half of ENSO is measured using the Southern Oscillation Index.

Timeframe: Every three months.

Region/Location: Niño3.4 region: 5°S – 5°N, 120° – 170°W

Measurement Platform: *In-situ* station, satellite, model.

Sourced from: NOAA Climate.gov and NOAA NCEI NCDC.




Figure 11. Oceanic Niño Index from 1950-2018 (top) and 2000–2018 (bottom) with El Niño periods in red and La Niña periods in blue

1.1.1.4 Pacific Decadal Oscillation (PDO)

Rationale: The Pacific Decadal Oscillation (PDO) was initially named by a fisheries scientist, Steven Hare, in 1996 while researching connections between Alaska salmon production cycles and Pacific climate. Like ENSO, the PDO reflects changes between periods of persistently warm or persistently cool ocean temperatures, but over a period of 20 to 30 years (versus 6 to 18 months for ENSO events). The climatic fingerprints of the PDO are most visible in the Northeastern Pacific, but secondary signatures exist in the tropics.

Status: The PDO was positive, or warm, for much of 2018. In March and June, the index dipped just below zero but returned to a positive value the following months. PDO index values are not yet available for the last three months of the year.




Figure 12. Pacific Decadal Oscillation from 1950–2018 (top) and 2000–2018 (bottom) with positive warm periods in red and negative cool periods in blue

Description: The Pacific Decadal Oscillation (PDO) is often described as a long-lived El Niño-like pattern of Pacific climate variability. As seen with the better-known El Niño – Southern Oscillation (ENSO), extremes in the PDO pattern are marked by widespread variations in the Pacific Basin and the North American climate. In parallel with the ENSO phenomenon, the extreme cases of the PDO have been classified as either warm or cool, as defined by ocean temperature anomalies in the northeast and tropical Pacific Ocean. When sea surface temperatures (SSTs) are below average in the interior North Pacific and warm along the North American coast, and when sea level pressures are below average in the North Pacific, the PDO has a positive value. When the climate patterns are reversed, with warm SST anomalies in the interior and cool SST anomalies along the North American coast, or above average sea level

pressures over the North Pacific, the PDO has a negative value. Description inserted from <https://www.ncdc.noaa.gov/teleconnections/pdo/>.

Timeframe: Annual, monthly

Region/Location: Pacific Basin north of 20°N.

Measurement Platform: *In-situ* station, satellite, model

Sourced from: NOAA ESRL (2019b).

1.1.1.5 Tropical Cyclones

Rationale: The effects of tropical cyclones are numerous and well known. At sea, storms disrupt and endanger shipping traffic as well as fishing effort and safety. The Hawai'i longline fishery, for example, has had serious problems with vessels dodging storms at sea, delayed departures, and inability to make it safely back to Honolulu because of bad weather. When cyclones encounter land, their intense rains and high winds can cause severe property damage, loss of life, soil erosion, and flooding. Associated storm surge, the large volume of ocean water pushed toward shore by cyclones' strong winds, can cause severe flooding and destruction.

Status:

Eastern North Pacific. Overall, the 2018 eastern Pacific hurricane season featured well above average activity. There were 22 named storms, of which 12 became hurricanes and 9 became major hurricanes - category 3 or higher on the Saffir-Simpson Hurricane Wind Scale. This compares to the long-term averages of 15 named storms, 8 hurricanes, and 4 major hurricanes. There were also 3 tropical depressions that did not reach tropical storm strength. In terms of Accumulated Cyclone Energy (ACE), which measures the strength and duration of tropical storms and hurricanes, activity in the basin in 2018 was the 3rd highest on record, behind 1990 and 1992. Summary inserted from <https://www.nhc.noaa.gov/text/MIATWSEP.shtml>.

Central North Pacific. Tropical cyclone activity in 2018 was high. The ACE index was the second highest since 1980, second only to 2015, and well above the 1981 – 2010 average of just under 20 ($\times 10^4$ knots²). Of note was Hurricane Lane, which reached Category 5 strength and passed within 110 miles of Honolulu. Lane was only the second Category 5 hurricane to pass within 250 miles of Hawaii, with the last being Hurricane John in 1994. Some of the impacts associated with Hurricane Lane include widespread reports of more than 40 inches of rain the islands of Hawaii and Kauai. There was one preliminary report of more than 52 inches of rain. At least one fatality was blamed on Hurricane Lane. Summary inserted from <https://www.ncdc.noaa.gov/sotc/tropical-cyclones/201808>.

Western North Pacific. Tropical cyclone activity was roughly average. The ACE Index was slightly above average in the Western North Pacific. Of note was Super Typhoon Yutu which made landfall on the islands of Tinian and Saipan as a Category 5 equivalent typhoon with estimated winds of 180 mph and a central minimum pressure of 905 mb. This marked the second strongest tropical cyclone to impact any U.S. territory on record. The storm devastated most of Tinian and Saipan with nearly every structure on the two islands being damaged or destroyed,

including the Saipan International Airport. There were two fatalities reported in the Northern Marianas. Summary inserted from <https://www.ncdc.noaa.gov/sotc/tropical-cyclones/201810>.

South Pacific. Tropical cyclone activity and the ACE Index were below average in 2018.

Description: This indicator uses historical data from the NOAA National Climate Data Center (NCDC) International Best Track Archive for Climate Stewardship to track the number of tropical cyclones in the western, central, eastern, and southern Pacific basins. This indicator also monitors the Accumulated Cyclone Energy (ACE) Index and the Power Dissipation Index which are two ways of monitoring the frequency, strength, and duration of tropical cyclones based on wind speed measurements.

The annual frequency of storms passing through each basin is tracked and a stacked time series plot shows the representative breakdown of Saffir-Simpson hurricane categories.

Every cyclone has an ACE Index value, which is a number based on the maximum wind speed measured at six-hourly intervals over the entire time that the cyclone is classified as at least a tropical storm (wind speed of at least 34 knots; 39 mph). Therefore, a storm's ACE Index value accounts for both strength and duration. This plot shows the historical ACE values for each hurricane/typhoon season and has a horizontal line representing the average annual ACE value.

Timeframe: Annual.

Region/Location:

Eastern North Pacific: east of 140° W, north of the equator.

Central North Pacific: 180° - 140° W, north of the equator.

Western North Pacific: west of 180°, north of the equator.

South Pacific: south of the equator.

Measurement Platform: Satellite.

Sourced from: NOAA NCEI (2018c)




Figure 13. 2018 Pacific basin tropical cyclone tracks




Figure 14. Tropical storm totals by region.




Figure 15. Accumulated Cyclone Energy (ACE) Index by region

1.1.1.6 Sea Surface Temperature (SST)

Rationale: Sea surface temperature is one of the most directly observable existing measures for tracking increasing ocean temperatures. SST varies in response to natural climate cycles such as the El Niño – Southern Oscillation (ENSO) and is projected to rise as a result of anthropogenic climate change. Both short-term variability and long-term trends in SST impact the marine ecosystem. Understanding the mechanisms through which organisms are impacted and the time scales of these impacts is an area of active research.

Status:

Pacific Remote Island Areas Grid: Annual mean SST was 28.22C in 2018. Over the period of record, monthly SST shows no significant pattern of increase or decrease. Monthly SST values in 2018 ranged from 26.60 – 29.22 °C, within the climatological range of 25.32 – 30.09 °C. The annual anomaly was 0.398 °C hotter than average, with mild intensification in the northern section of the grid.

Johnston Atoll Grid: Annual mean SST was 26.73°C in 2018. Over the period of record, annual SST has increased at a rate of 0.018 °C yr⁻¹. Monthly SST values in 2018 ranged from 25.48 – 28.48°C, within the climatological range of 24.37 – 28.79 °C. The annual anomaly was 0.344°C hotter than average, with no dramatic spatial pattern.

Wake Atoll Grid: Annual mean SST was 27.44°C in 2018. Over the period of record, annual SST has increased at a rate of 0.024 °C yr⁻¹. Monthly SST values in 2018 ranged from 25.82 – 29.27°C, within the climatological range of 24.23 – 29.97 °C. The annual anomaly was 0.153°C hotter than average, with no dramatic spatial pattern.

Note that from the top to bottom in Figure 16, panels show climatological SST (1982-2017), 2018 SST anomaly, time series of monthly mean SST, and time series of monthly SST anomaly. The white box in the upper panels indicates the area over which SST is averaged for the time series plots.


Description: Satellite remotely-sensed monthly sea surface temperature (SST) is averaged across each of the PRIA Grid (1°S – 7°N, 159° – 177°W; including Howland, Baker, Jarvis, Palmyra, Kingman Reef), Johnston Island (16° – 17°N, 168° – 170°W), and Wake Atoll (17.7° – 20.7°N, 165° – 168°W). Time series of monthly mean SST averaged over the respective grids are presented. Additionally, spatial climatology and anomalies are shown. Data from NOAA Pathfinder v5.3.

Timeframe: Monthly.


Region/Location: PRIA Grid (1°S – 7°N, 159° – 177°W); Johnston Atoll (16° – 17°N, 168° – 170°W), and Wake Atoll (17.7° – 20.7°N, 165° – 168°W)

Measurement Platform: Satellite.


Sourced from: NOAA OceanWatch (2018).



1982-2017
Climatology



2018
Anomaly




Monthly Mean
1982 - 2017
2018




Monthly Anomaly
1982 - 2017
2018


Figure 16. Sea surface temperature climatology and anomalies from the PRIA Grid




1982-2017
Climatology



2018
Anomaly




Monthly Mean
1982 - 2017
2018




Monthly Anomaly
1982 - 2017
2018


Figure 17. Sea surface temperature climatology and anomalies from Johnston Atoll Grid




1982-2017
Climatology



2018
Anomaly



Monthly Mean
1982 - 2017
2018



Monthly
Anomaly
1982 - 2017
2018

Figure 18. Sea surface temperature climatology and anomalies from Wake Atoll Grid

2.4.3.7 Coral Thermal Stress Exposure: Degree Heating Weeks

Rationale: Degree heating weeks are one of the most widely used metrics for assessing exposure to coral bleaching-relevant thermal stress.

Description: Here we present a metric of exposure to thermal stress that is relevant to coral bleaching. Degree Heating Weeks (DHW) measure time and temperature above a reference ‘summer maximum’, presented as a rolling sum weekly thermal anomalies over a 12-week window. Higher DHW measures imply a greater likelihood of mass coral bleaching or mortality from thermal stress.

Status: After a major heat stress events in 2015- 2016 in all three virtual stations, the Northern line island showed a minor event in 2018.

The NOAA Coral Reef Watch program uses satellite data to provide current reef environmental conditions to quickly identify areas at risk for [coral bleaching](#). Bleaching is the process by which corals lose the symbiotic algae that give them their distinctive colors. If a coral is severely bleached, disease and death become likely.

The NOAA Coral Reef Watch (CRW) daily 5-km satellite coral bleaching Degree Heating Week (DHW) product presented here shows accumulated heat stress, which can lead to coral bleaching and death. The scale goes from 0 to 20 °C-weeks. The DHW product accumulates the instantaneous bleaching heat stress (measured by Coral Bleaching HotSpots) during the most-recent 12-week period. It is directly related to the timing and intensity of coral bleaching. Significant coral bleaching usually occurs when DHW values reach 4 °C-weeks. By the time DHW values reach 8 °C-weeks, widespread bleaching is likely and significant mortality can be expected.

Timeframe: 2013-2018, Daily data.

Region/Location: Global.

Data Source: “NOAA Coral Reef Watch” <https://coralreefwatch.noaa.gov>

Measurement Platform: [NOAA/NESDIS operational daily global 5km geostationary-polar-orbiting \(Geo-Polar\) Blended Night-only SST Analysis](#)




Figure 19. Coral Thermal Stress Exposure, Howland/Baker Virtual Station 2013-2018 (Coral Reef Watch Degree Heating Weeks)




Figure 20. Coral Thermal Stress Exposure, Northern Line Islands Virtual Station 2013-2018 (Coral Reef Watch Degree Heating Weeks)




Figure 21. Coral Thermal Stress Exposure, Johnston Virtual Station 2013-2018 (Coral Reef Watch Degree Heating Weeks)




Figure 22. Coral Thermal Stress Exposure, Wake Atoll Virtual Station 2013-2018 (Coral Reef Watch Degree Heating Weeks)

2.4.3.8 Chlorophyll-A and Anomaly

Rationale: Chlorophyll-A is one of the most directly observable measures we have for tracking increasing ocean productivity.

Status:

Pacific Remote Island Areas: Annual mean Chl-A was 0.144 mg/m³ in 2018. Over the period of record, annual Chl-A has shown a significant linear decrease at a rate of 0.00138 mg/m³. Monthly Chl-A values in 2018 ranged from 0.122-0.164 mg/m³, within the climatological range of 0.049 – 0.224 mg/m³. The annual anomaly was 0.013 mg/m³ lower than climatological values, with lower values along the equator, and higher values toward the west and poleward.

Johnston Atoll: Annual mean Chl-A was 0.055 mg/m³ in 2018. Over the period of record, annual Chl-A has shown no significant linear change. Monthly Chl-A values in 2018 ranged from 0.049-0.062 mg/m³, within the climatological range of 0.036 – 0.090 mg/m³. The annual anomaly was roughly in-line with climatological values (0.001 mg/m³ lower), with an intensification toward the west of the atoll.

Wake Atoll: Annual mean Chl-A was 0.052 mg/m³ in 2018. Over the period of record, annual Chl-A has shown no significant linear change. Monthly Chl-A values in 2018 ranged from 0.048-0.057 mg/m³, within the climatological range of 0.035 – 0.114 mg/m³. The annual anomaly was roughly in-line with climatological values (0.000081 mg/m³ higher), with an intensification toward the west of the atoll.

Description: Chlorophyll-A Concentration from July 1998- June 2018 (data for the 2d half of 2018 is not yet available), derived from the ESA Ocean Color Climate Change Initiative dataset, a multi-sensor global dataset. A monthly climatology was generated across the entire period (1998-2018) to provide both a 2018 spatial anomaly, and an anomaly time series.

The Ocean Colour Climate Change Initiative project aims to:

- Develop and validate algorithms to meet the Ocean Colour GCOS ECV requirements for consistent, stable, error-characterized global satellite data products from multi-sensor data archives.
- Produce and validate, within an R&D context, the most complete and consistent possible time series of multi-sensor global satellite data products for climate research and modelling.
- Optimize the impact of MERIS data on climate data records.
- Generate complete specifications for an operational production system.
- Strengthen inter-disciplinary cooperation between international Earth observation, climate research and modelling communities, in pursuit of scientific excellence.

The ESA OC CCI project is following a data reprocessing paradigm of regular re-processing utilizing on-going research and developments in atmospheric correction, in-water algorithms, data merging techniques and bias correction. This requires flexibility and rapid turn-around of processing of extensive ocean color datasets from a number of ESA and NASA missions to both trial new algorithms and methods and undertake the complete data set production.


Data products generated by the Ocean Colour component of the European Space Agency Climate Change Initiative project. These files are daily composites of merged sensor (MERIS, Moderate Resolution Imaging Spectroradiometer (MODIS) Aqua, Sea-Wide Field-of-View Sensor (SeaWiFS) Local Area Coverage (LAC) & Global Area Coverage (GAC), Visible and Infrared Imager/Radiometer Suite (VIIRS)) products. MODIS Aqua and MERIS were band-shifted and bias-corrected to SeaWiFS bands and values using a temporally and spatially varying scheme based on the overlap years of 2003-2007. VIIRS was band-shifted and bias-corrected in a second stage against the MODIS Rrs that had already been corrected to SeaWiFS levels, for the overlap period 2012-2013. VIIRS and SeaWiFS Rrs were derived from standard NASA L2 products; MERIS and MODIS from a combination of NASA's l2gen (for basic sensor geometry corrections, etc.) and HYGEOS Polymer v3.5 (for atmospheric correction). The Rrs were binned to a sinusoidal 4km level-3 grid, and later to 4km geographic projection, by Brockmann Consult's BEAM. Derived products were generally computed with the standard SeaDAS algorithms. QAA IOPs were derived using the standard SeaDAS algorithm but with a modified backscattering table to match that used in the bandshifting. The final chlorophyll is a combination of OC4, Hu's CI and OC5, depending on the water class memberships. Uncertainty estimates were added using the fuzzy water classifier and uncertainty estimation algorithm of Tim Moore as documented in Jackson et al (2017).

Timeframe: July 1998 – June 2018, Daily data available, Monthly means shown.


Region/Location: Global.

Data Source: “ESA OC CCI (ERDDAP Monthly)” <http://oceanwatch.pifsc.noaa.gov/doc.html>


Measurement Platform: *Multi-Sensor Dataset*.




1998-2017
Climatology



2018
Anomaly




Monthly Mean
1998 - 2017
2018




Monthly Anomaly
1998 - 2017
2018


Figure 23. Chlorophyll-A (Chl-A) and Chl-A Anomaly from the PRIA Grid



1998-2017
Climatology



2018
Anomaly




Monthly Mean
1998 - 2017
2018




Monthly Anomaly
1998 - 2017
2018


Figure 24. Chlorophyll-A (Chl-A) and Chl-A Anomaly from the Johnston Atoll Grid




1998-2017
Climatology



2018
Anomaly



Monthly Mean
1998 - 2017
2018



Monthly
Anomaly
1998 - 2017
2018

Figure 25. Chlorophyll-A (Chl-A) and Chl-A Anomaly from the Wake Atoll Grid

2.4.3.9 Rainfall (CMAP Precipitation)

Rationale: Rainfall may have substantive effects on the nearshore environment and is a potentially important co-variate with the landings of particular stocks.

Description: The CPC Merged Analysis of Precipitation ("CMAP") is a technique which produces pentad and monthly analyses of global precipitation in which observations from rain gauges are merged with precipitation estimates from several satellite-based algorithms (infrared and microwave). The analyses are on a 2.5 x 2.5 degree latitude/longitude grid and extend back to 1979. CMAP Precipitation data provided by the NOAA/OAR/ESRL PSD, Boulder, Colorado, USA, from their Web site at <https://www.esrl.noaa.gov/psd/>. These data are comparable (but should not be confused with) similarly combined analyses by the [Global Precipitation Climatology Project](#) which are described in Huffman et al (1997).

It is important to note that the input data sources to make these analyses are not constant throughout the period of record. For example, SSM/I (passive microwave - scattering and emission) data became available in July of 1987; prior to that the only microwave-derived estimates available are from the MSU algorithm (Spencer1993) which is emission-based thus precipitation estimates are available only over oceanic areas. Furthermore, high temporal resolution IR data from geostationary satellites (every 3-hr) became available during 1986; prior to that, estimates from the OPI technique (Xie and Arkin 1997) are used based on OLR from polar orbiting satellites.

The merging technique is thoroughly described in Xie and Arkin (1997). Briefly, the methodology is a two-step process. First, the random error is reduced by linearly combining the satellite estimates using the maximum likelihood method, in which case the linear combination coefficients are inversely proportional to the square of the local random error of the individual data sources. Over global land areas the random error is defined for each time period and grid location by comparing the data source with the rain gauge analysis over the surrounding area. Over oceans, the random error is defined by comparing the data sources with the rain gauge observations over the Pacific atolls. Bias is reduced when the data sources are blended in the second step using the blending technique of Reynolds (1988). Here the data output from step 1 is used to define the "shape" of the precipitation field and the rain gauge data are used to constrain the amplitude.

Timeframe: Monthly

Region/Location: Global

Measurement Platform: *In-situ* station gauges and satellite data.




Figure 26. CMAP precipitation across the PRIA Grid with 2018 values in red




Figure 27. CMAP precipitation across the Johnston Atoll Grid with 2018 values in red




Figure 28. CMAP precipitation across the Wake Atoll Grid with 2018 values in red

2.4.3.10 Sea Level (Sea Surface Height and Anomaly)

Rationale: Rising coastal sea levels can result in a number of coastal impacts, including inundation of infrastructure, increased damage resulting from storm-driven waves and flooding, and saltwater intrusion into freshwater supplies.

Description: Monthly mean sea level time series, including extremes

Timeframe: Monthly

Region/Location: Observations from selected sites within the Marianas Archipelago

Measurement Platform: Satellite and *in situ* tide gauges

2.4.3.10.1 Basin-Wide Perspective

This image of the mean sea level anomaly for March 2019 compared to 1993-2013 climatology from satellite altimetry provides a glimpse into how the current weak El Niño continues to affect sea level across the Pacific Basin. The image captures the fact that sea level continues to be lower in the Western Pacific and higher in the Central and Eastern Pacific (a standard pattern during El Niño events. This basin-wide perspective provides a context for the location-specific sea level/sea surface height images that follow.)




Figure 29a. Sea surface height and anomaly across the Pacific Ocean





Figure 29b. Quarterly time series of mean sea level anomalies during 2018 show no pattern of El Niño throughout the year according to satellite altimetry measurements of sea level height (unlike 2015).



2.4.3.10.2 Local Sea Level

These time-series from *in situ* tide gauges provide a perspective on sea level trends within each Archipelago (Tide Station Time Series from NOAA/COOPS).

The following figures and descriptive paragraphs were inserted from NOAA (2018). Figure 30 and Figure 31 show the monthly mean sea level without the regular seasonal fluctuations due to coastal ocean temperatures, salinities, winds, atmospheric pressures, and ocean currents. The long-term linear trend is also shown, including its 95% confidence interval. The plotted values are relative to the most recent [Mean Sea Level datum established by CO-OPS](#). The calculated trends for all stations are available as a [table in millimeters/year and in feet/century](#) (0.3 meters = 1 foot). If present, solid vertical lines indicate times of any major earthquakes in the vicinity of the station and dashed vertical lines bracket any periods of questionable data or datum shift.

The relative sea level trend is 0.75 millimeters/year with a 95% confidence interval of +/- 0.56 mm/yr based on monthly mean sea level data from 1947 to 2003 which is equivalent to a change of 0.25 feet in 100 years (Figure 30).

The relative sea level trend is 2.1 millimeters/year with a 95% confidence interval of +/- 0.42 mm/yr based on monthly mean sea level data from 1950 to 2018 which is equivalent to a change of 0.69 feet in 100 years (Figure 31).




Figure 30. Monthly mean sea level without the regular seasonal fluctuations due to coastal ocean temperatures, salinities, winds, atmospheric pressures, and ocean currents




Figure 31. Monthly mean sea level without the regular seasonal fluctuations due to coastal ocean temperatures, salinities, winds, atmospheric pressures, and ocean currents

2.4.3.1 References

- Aviso, 2019. El Niño Bulliten – Latest plots. Ocean indicator products, Center for Topographic Studies of the Ocean and Hydrosphere, Centre National D'études Spatiales. Available at <https://www.aviso.altimetry.fr/en/data/products/ocean-indicators-products/el-nino-bulletin.html>. Updated April 2019.
- Fabry, V.J., Seibel, B.A., Feely, R.A., and J.C. Orr, 2008. Impacts of ocean acidification on marine fauna and ecosystem processes. *ICES Journal of Marine Science*, 65, pp. 414-432.
- Feely, R.A., Alin, S.R., Carter, B., Bednarsek, N., Hales, B., Chan, F., Hill, T.M., Gaylord, B., Sanford, E., Byrne, R.H., Sabine, C.L., Greeley, D., and L. Juranek, 2016. Chemical and biological impacts of ocean acidification along the west coast of North America. *Estuarine, Coastal and Shelf Science*, 183, pp. 260-270. doi: 10.1016/j.ecss.2016.08.043.
- Huffman, G.J., Adler, R.F., Arkin, P., Chang, A., Ferraro, R., Gruber, A., Janowiak, J., McNab, A., Rudolf, B. and U. Schneider, 1997. The global precipitation climatology project (GPCP) combined precipitation dataset. *Bulletin of the American Meteorological Society*, 78(1), pp.5-20.
- Kanamitsu, M., Ebisuzaki, W., Woollen, J., Yang, S.K., Hnilo, J.J., Fiorino, M. and G.L. Potter, 2002. NCEP–DOE AMIP-II Reanalysis (R-2), *B. Am. Meteorol. Soc.*, 83, pp. 1631–1643,
- Keeling, C.D., Bacastow, R.B., Bainbridge, A.E., Ekdahl, C.A., Guenther, P.R., and L.S. Waterman, 1976. Atmospheric carbon dioxide variations at Mauna Loa Observatory, Hawaii. *Tellus*, 28, pp. 538-551.
- Knapp, K.R., M.C. Kruk, D.H. Levinson, H.J. Diamond, and C.J. Neumann, 2010: The International Best Track Archive for Climate Stewardship (IBTrACS): Unifying

- tropical cyclone best track data. *Bulletin of the American Meteorological Society*, 91, pp. 363-376. doi:10.1175/2009BAMS2755.1.
- Li, X., Pichel, W.G., Clemente-Colón, P., and J.F. Sapper, 2001a. Deriving the operational nonlinear multi-channel sea surface temperature algorithm coefficients for NOAA-15 AVHRR/3, *Int. J. Remote Sens.*, 22(4), pp. 699 - 704.
- Li, X., Pichel, W.G., Clemente-Colón, P., Krasnopolsky, V., and J.F Sapper, 2001b. Validation of coastal sea and lake surface temperature measurements derived from NOAA/AVHRR Data, *Int. J. Remote Sens.*, 22(7), pp. 1285-1303.
- Liu, G., Heron, S.F., Eakin, C.M., Muller-Karger, F.E., Vega-Rodriguez, M., Guild, L.S., De La Cour, J.L., Geiger, E.F., Skirving, W.J., Burgess, T.F. and A.E. Strong, 2014. Reef-scale thermal stress monitoring of coral ecosystems: new 5-km global products from NOAA Coral Reef Watch. *Remote Sensing*, 6(11), pp.11579-11606.
- Mantua, N., 2018. The Pacific Decadal Oscillation. Available at <http://research.jisao.washington.edu/pdo/>. Accessed March 2018.
- Mélin, F., V. Vantrepotte, A. Chuprin, M. Grant, T. Jackson, and S. Sathyendranath, 2017. "Assessing the fitness-for-purpose of satellite multi-mission ocean color climate data records: A protocol applied to OC-CCI chlorophyll-a data." *Remote sensing of environment*, 203, 139-151.
- NOAA, 2002. CPC Merged Analysis of Precipitation. National Weather Service, National Centers for Environmental Prediction, Climate Prediction Center. Available at https://www.cpc.ncep.noaa.gov/products/global_precip/html/wpage.cmap.html. Updated 25 September 2002.
- NOAA, 2018. Sea level Trends. NOAA Tides & Currents. Available at https://tidesandcurrents.noaa.gov/sltrends/sltrends_station.shtml?stnid=1770000. Updated 8 August 2018.
- NOAA, 2019a. Chlorophyll a Concentration, Aqua MODIS. OceanWatch – Central Pacific Node. NASA/GSFC/OBPG. Available at <https://oceanwatch.pifsc.noaa.gov/contact.html>. Updated 21 February 2019.
- NOAA, 2019b. Full Mauna Loa CO₂ record. Trends in Atmospheric Carbon Dioxide. NOAA Earth System Research Laboratory, Global Monitoring Division. Available from <https://www.esrl.noaa.gov/gmd/ccgg/trends/full.html>. Updated 28 May 2019.
- NOAA, 2019c. Sea Surface Temperature Datasets. AVHRR Pathfinder v.5, AVRR GAC v.5, and GOES-POES v.5. OceanWatch – Central Pacific Node. NASA/GSFC/OBPG. Available at <https://oceanwatch.pifsc.noaa.gov/contact.html>. Updated 21 February 2019.
- NOAA Coral Reef Watch, 2019, updated daily. Version 3.1 Daily Global 5-km Satellite Coral Bleaching Degree Heating Week Product, 2013-2018. College Park, Maryland, USA: NOAA Coral Reef Watch. Available at <https://coralreefwatch.noaa.gov/satellite/hdf/index.php>.

- NOAA NCEI, 2019. State of the Climate: Hurricanes and Tropical Storms for Annual 2018. Available from <http://www.ncdc.noaa.gov/sotc/tropical-cyclones/201713>. Accessed 30 March 30, 2018.
- NOAA PMEL, 2019. A primer on pH. NOAA Pacific Marine Environmental Laboratory, Carbon Program. Available at <https://www.pmel.noaa.gov/co2/story/A+primer+on+pH>.
- Reynolds, R.W., 1988. A real-time global sea surface temperature analysis. *Journal of Climate*, 1(1), pp. 75-87.
- Savchenko, A., Ouzounov, D., Ahmad, S., Acker, J., Leptoukh, G., Koziana, J., and D. Nickless, 2004. Terra and Aqua MODI products available from NASA GES DAAC. *Advances in Space Research*, 34(4), pp. 710-714.
- Spencer, R.W., 1993. Global oceanic precipitation from the MSU during 1979—91 and comparisons to other climatology. *Journal of Climate*, 6(7), pp.1301-1326.
- State of Environmental Conditions in Hawaii and the U.S. Affiliated Pacific Islands under a Changing Climate, 2017. Coordinating Authors: J.J. Marra and M.C. Kruk. Contributing Authors: M. Abecassis; H. Diamond; A. Genz; S.F. Heron; M. Lander; G. Liu; J. T. Potemra; W.V. Sweet; P. Thompson; M.W. Widlansky; and P. Woodworth-Jefcoats. September, 2017. NOAA NCEI.
- Stowe, L.L., Davis, P.A., and E.P. McClain, 1999. Scientific basis and initial evaluation of the CLAVR-1 global clear/cloud classification algorithm for the advanced very high resolution radiometer. *J. Atmos. Oceanic Technol.*, 16, pp. 656-681.
- Thoning, K.W., Tans, P.P., and W.D. Komhyr, 1989. Atmospheric carbon dioxide at Mauna Loa Observatory 2. Analysis of the NOAA GMCC data, 1974-1985. *Journal of Geophysical Research*, 94, pp. 8549-8565.
- Walton C.C., Pichel, W.G., Sapper, J.F., and D.A. May, 1998. The development and operational application of nonlinear algorithms for the measurement of sea surface temperatures with the NOAA polar-orbiting environmental satellites, *J. Geophys. Res.*, 103(C12), pp. 27999-28012.
- Xie, P. and P.A. Arkin, 1997. Global precipitation: A 17-year monthly analysis based on gauge observations, satellite estimates, and numerical model outputs. *Bulletin of the American Meteorological Society*, 78(11), pp. 2539-2558.

2.5 ESSENTIAL FISH HABITAT

2.5.1 Introduction

The Magnuson-Stevens Fishery Conservation and Management Act (MSA) includes provisions concerning the identification and conservation of essential fish habitat (EFH), and under the EFH final rule, habitat areas of particular concern (HAPC) (50 Code of Federal Regulations [CFR] 600.815). The MSA defines EFH as “those waters and substrate necessary to fish for spawning, breeding, feeding, or growth to maturity.” HAPC are those areas of EFH identified pursuant to 50 CFR 600.815(a)(8), and meeting one or more of the following considerations: (1) ecological function provided by the habitat is important; (2) habitat is sensitive to human-induced environmental degradation; (3) development activities are, or will be, stressing the habitat type; or (4) the habitat type is rare.

NMFS and the regional Fishery Management Councils must describe and identify EFH in fishery management plans (FMPs), minimize to the extent practicable the adverse effects of fishing on EFH, and identify other actions to encourage the conservation and enhancement of EFH. Federal agencies that authorize, fund, or undertake actions that may adversely affect EFH must consult with NMFS, and NMFS must provide conservation recommendations to federal and state agencies regarding actions that would adversely affect EFH. Councils also have the authority to comment on federal or state agency actions that would adversely affect the habitat, including EFH, of managed species.

The EFH Final Rule strongly recommends regional fisheries management councils and NMFS to conduct a review and revision of the EFH components of fisheries management plans every five years (600.815(a)(10)). The council’s FEPs state that new EFH information should be reviewed, as necessary, during preparation of the annual reports by the Plan Teams. Additionally, the EFH Final Rule states “Councils should report on their review of EFH information as part of the annual Stock Assessment and Fishery Evaluation (SAFE) report prepared pursuant to §600.315(e).” The habitat portion of the annual SAFE report is designed to meet the FEP requirements and EFH Final Rule guidelines regarding EFH reviews.

National Standard 2 guidelines recommend that the SAFE report summarize the best scientific information available concerning the past, present, and possible future condition of EFH described by the FEPs.

2.5.2 EFH Information

The EFH components of fisheries management plans include the description and identification of EFH, lists of prey species and locations for each managed species, and optionally, habitat areas of particular concern. Impact-oriented components of FMPs include federal fishing activities that may adversely affect EFH; non-federal fishing activities that may adversely affect EFH; non-fishing activities that may adversely affect EFH; conservation and enhancement recommendations; and a cumulative impacts analysis on EFH. The last two components include the research and information needs section, which feeds into the Council’s Five Year Research Priorities, and the EFH update procedure, which is described in the FEP but implemented in the annual SAFE report.

The Council has described EFH for five management unit species (MUS) under its management authority: pelagic (PMUS), bottomfish (BMUS), crustaceans (CMUS), coral reef ecosystem (CREMUS), and precious corals (PCMUS). The PRIA FEP describes EFH for the BMUS, CMUS, CREMUS, and PCMUS.

EFH reviews of the biological components, including the description and identification of EFH, lists of prey species and locations, and HAPC, consist of three to four parts:

- Updated species descriptions, which can be found appended to previous SAFE reports and can be used to directly update the FEP;
- Updated EFH levels of information tables, which can be found in Section 2.5.5;
- Updated research and information needs, which can be found in Section 2.5.6 and can be used to directly update the FEP; and
- An analysis that distinguishes EFH from all potential habitats used by the species, which is the basis for an options paper for the Council and can be developed if enough information exists to refine EFH.

2.5.2.1 Habitat Objectives of FEP

The habitat objective of the FEP is to refine EFH and minimize impacts to EFH, with the following sub-objectives:

- Review EFH and HAPC designations every five years based on the best available scientific information and update such designations based on the best available scientific information, when available
- Identify and prioritize research to: assess adverse impacts to EFH and HAPC from fishing (including aquaculture) and non-fishing activities, including, but not limited to, activities that introduce land-based pollution into the marine environment.

The annual reports have reviewed the precious coral EFH components and non-fishing impacts components. The Council's support of non-fishing activities research is monitored through the program plan and five year research priorities, not the annual report.

2.5.2.2 Response to Previous Council Recommendations

At its 172nd meeting in March 2018, the Council recommended that staff develop an omnibus amendment updating the non-fishing impact to EFH sections of the FEPs, incorporating the non-fishing impacts EFH review report by Minton (2017) by reference. An options paper has been developed.

At its 173rd meeting in June 2018, the Council directed staff to develop options to redefine EFH and any HAPC for precious corals in Hawaii for Council consideration for an FEP amendment. An options paper was developed.

At its 174th meeting in October 2018, the Council directed staff to prepare an amendment to the Hawaii FEP to revise the Precious Corals EFH and selected the following preliminarily preferred options for the staff to further analyze:

- Action 1: Option 4 - Revise existing beds and designate new beds as EFH
- Action 2: Option 2 - Update Geographic Extent and Habitat Characteristics
- Action 3: Option 1 - Update the FEPs

An FEP amendment is being developed to present to the Council in 2019.

2.5.3 Habitat Use by MUS and Trends in Habitat Condition

The PRIAs comprise the U.S. possessions of Baker Island, Howland Island, Jarvis Island, Johnston Atoll, Kingman Reef, Wake Island, Palmyra Atoll, and Midway Atoll (Figure 32). However, because Midway is located in the Hawaiian archipelago, it is included in the Hawaii Archipelago FEP¹. Therefore, neither the “Pacific Remote Island Areas” nor “PRIA” include Midway Atoll, for the purpose of federal fisheries management.

Baker Island is part of the Phoenix Islands archipelago. It is located approximately 1,600 nautical miles (nm) to the southwest of Honolulu at 0° 13' N and 176° 38' W. Baker is a coral-topped seamount surrounded by a narrow-fringing reef that drops steeply very close to the shore. The total amount of emergent land area of Baker Island is 1.4 square kilometers.

Howland Island lies approximately 35 miles due north of Baker Island and is also part of the Phoenix Islands archipelago. The island, which is the emergent top of a seamount, is fringed by a relatively flat coral reef that drops off sharply. Howland Island is approximately 1.5 miles long and 0.5 miles wide. The island is flat and supports some grasses and small shrubs. The total land area is 1.6 square kilometers.

Jarvis Island, which is part of the Line Island archipelago, is located approximately 1,300 miles south of Honolulu and 1,000 miles east of Baker Island. It sits 23 miles south of the Equator at 160° 01' W. Jarvis Island is a relatively flat, sandy coral island with a 15–20-ft beach rise. Its total land area is 4.5 square kilometers. It experiences a very dry climate.

Palmyra Atoll is a low-lying coral atoll system comprised of approximately 52 islets surrounding three central lagoons. It is approximately 1,050 nm south of Honolulu and is located at 5° 53' N and 162° 05' W. It is situated about halfway between Hawaii and American Samoa. Palmyra Atoll is located in the intertropical convergence zone, an area of high rainfall.

Kingman Reef is located 33 nm northwest of Palmyra Atoll at 6° 23' N and 162° 24' W. Along with Palmyra, it is at the northern end of the Line Island archipelago. Kingman is actually a series of fringing reefs around a central lagoon with no emergent islets that support vegetation.

Wake Island is located at 19° 18' N and 166° 35' E, and is the northernmost atoll of the Marshall Islands group, located approximately 2,100 miles west of Hawaii. Wake Island has a total land area of 6.5 square kilometers and comprises three islets: Wake, Peale, and Wilkes.

¹ Midway is not administered civilly by the State of Hawaii.

Johnston Atoll is located at 16° 44' N and 169° 31' W and is approximately 720 nm southwest of Honolulu. French Frigate Shoals in the NWHI, about 450 nm to the northwest, is the nearest land mass. Johnston Atoll is an egg-shaped coral reef and lagoon complex comprised of four small islands totaling 2.8 square kilometers. The complex resides on a relatively flat, shallow platform approximately 34 kilometers in circumference. Johnston Island, the largest and main island, is natural, but has been enlarged by dredge-and-fill operations. Sand Island is composed of a naturally-formed island on its eastern portion and is connected by a narrow, man-made causeway to a dredged coral island at its western portion. The remaining two islands, North Island and East Island, are completely man-made from dredged coral.

All commercial activity is prohibited within the Pacific Remote Island Area Marine National Monument, which is 50 nm surrounding Palmyra Atoll and Kingman Reef and Howland and Baker Islands, and the entire US EEZ surrounding Johnston Atoll, Wake, and Jarvis Island.

Essential fish habitat in the PRIA for the four MUS comprises all substrate from the shoreline to the 700 m isobath (Figure 33). The entire water column is described as EFH from the shoreline to the 700 m isobath, and the water column to a depth of 400 m is described as EFH from the 700 m isobath to the limit or boundary of the exclusive economic zone (EEZ). While the coral reef ecosystems surrounding the islands in the PRIA have been the subject of a comprehensive monitoring program through the PIFSC Coral Reef Ecosystem Division (CRED) biennially since 2002, surveys are focused on the nearshore environments surrounding the islands, atolls, and reefs (PIBHMC).

The mission of the PIFSC Coral Reef Ecosystem Division (CRED) is to “provide high-quality, scientific information about the status of coral reef ecosystems of the U.S. Pacific islands to the public, resource managers, and policymakers on local, regional, national, and international levels” (PIFSC, 2011). CRED’s Reef Assessment and Monitoring Program (RAMP) conducts comprehensive ecosystem monitoring surveys at about 50 islands, atolls, and shallow bank sites in the Western Pacific Region on a one to three year schedule (Brainard et al., 2006). CRED coral reef monitoring reports provide the most comprehensive description of nearshore habitat quality in the region. The benthic habitat mapping program provides information on the quantity of habitat.




Figure 32. Pacific Remote Island Areas and the associated Pacific Remote Islands Marine National Monument




Figure 33. The Substrate EFH Limit and 700 meter isobath around the PRIAs (from Ryan et al. 2009)

2.5.3.1 Habitat Mapping

Mapping products for the PRIA are available from the Pacific Islands Benthic Habitat Mapping Center, and are listed in Table 5.

Table 5. Summary of habitat mapping in the PRIA

Depth Range	Timeline/Mapping Product	Progress	Source
0-30 m	IKONOS Benthic Habitat Maps	Palmyra only	Miller et al. (2011)
	2000-2010 Bathymetry	67%	DesRochers (2016)
	2011-2015 Multibeam Bathymetry		DesRochers (2016)
	2011-2015 Satellite Worldview 2 Bathymetry	Wake, Baker, and Howland Islands, Johnston and Palmyra Atolls, and Kingman Reef	Pers. Comm. DesRochers, March 19, 2018
30-150 m	2000-2010 Bathymetry	79%	DesRochers (2016)
	2011-2015 Multibeam Bathymetry	Howland and Baker updated with data collected in a few small areas in 2015	Pers. Comm., DesRochers, March 19, 2018
15 to 2500 m	Multibeam bathymetry	Complete at Jarvis, Howland, and Baker Islands	Pacific Islands Benthic Habitat Mapping Center
	Derived Products	Backscatter available for all Geomorphology products for Johnston, Howland, Baker, Wake	Pacific Islands Benthic Habitat Mapping Center

The land and seafloor area surrounding the islands and atolls of the PRIA are reproduced from Miller et al. (2011) and shown in Figure 34 alongside other physical data.




Figure 34. PRIA Land and Seafloor Primary Data Coverage (from Miller et al., 2011)

2.5.3.2 Benthic Habitat

Juvenile and adult life stages of coral reef MUS and crustaceans including spiny and slipper lobsters and Kona crab extends from the shoreline to the 100 m isobath (64 FR 19067, 19 April 1999). All benthic habitat is considered EFH for crustaceans species (64 FR 19067, 19 April 1999), while the type of bottom habitat varies by family for coral reef species (69 FR 8336, 24 February 2004). Juvenile and adult bottomfish EFH extends from the shoreline to the 400 m isobath (64 FR 19067, 19 April 1999), and juvenile and adult deepwater shrimp habitat extends from the 300 m isobath to the 700 m isobath (73 FR 70603, 21 November 2008). Table 6 shows the depths of geologic features, the occurrence of MUS EFH at that feature, and the availability of long-term monitoring data at diving depths.

Table 6. Occurrence of EFH by feature in the PRIAs

Feature	Summit Minimum Depth	Coral Reef/Crustaceans (w/o Deepwater Shrimp)	Bottomfish	Deepwater Shrimp	CRED Long Term Monitoring
Johnston Atoll	Emergent	✓	✓	✓	✓
Palmyra	Emergent	✓	✓	✓	✓
Kingman Reef	Emergent	✓	✓	✓	✓
Extensive banks 80 km SW of Kingman		?	?	?	
Jarvis Island	Emergent	✓	✓	✓	✓
Howland Island	Emergent	✓	✓	✓	✓
Baker Island	Emergent	✓	✓	✓	✓
Southeast of Baker	?	?	?	✓	
Wake Island	Emergent	✓	✓	✓	✓
South of Wake	?	?	?	✓	

2.5.3.2.1 RAMP Indicators

Benthic percent cover of coral, macroalgae, and crustose coralline algae from CRED are found in the following tables. CRED uses the benthic towed-diver survey method to monitor changes in benthic composition. In this method, “a pair of scuba divers (one collecting fish data, the other collecting benthic data) is towed about one m above the reef roughly 60 m behind a small boat at a constant speed of about 1.5 kt. Each diver maneuvers a tow board platform, which is connected

to the boat by a bridle and towline, and is outfitted with a communications telegraph and various survey equipment, including a downward-facing digital SLR camera (Canon EOS 50D, Canon Inc., Tokyo). The benthic towed diver records general habitat complexity and type (e.g., spur and groove, pavement), percent cover by functional-group (hard corals, stressed corals, soft corals, macroalgae, crustose coralline algae, sand, and rubble) and for macroinvertebrates (crown-of-thorns sea stars, sea cucumbers, free and boring urchins, and giant clams).

Towed-diver surveys are typically 50 min long and cover about two to three km of habitat. Each survey is divided into five-minute segments, with data recorded separately per segment to allow for later location of observations within the ~ 200-300 m length of each segment. Throughout each survey, latitude and longitude of the survey track are recorded on the small boat using a GPS; and after the survey, diver tracks are generated with the GPS data and a layback algorithm that accounts for position of the diver relative to the boat” (McCoy et al. 2017). The most recent data collected were in 2016 and described by McCoy et al. (2017).

Table 7. Mean percent cover of live coral from RAMP sites collected from towed-diver surveys in the PRIA

Year	2001	2002	2004	2005	2006	2007	2008	2009	2010	2011	2012	2014	2015	2016
Baker	35.37	49.47	38.78		32.95		41.20		47.44		42.10		34.48	
Howland	29.06	42.53	36.75		34.69		44.47		50.74		43.26		23.20	
Jarvis	24.22	26.19	30.63		28.54		27.70		26.92		25.38		39.75	
Johnston			5.01		22.95		18.38		7.94		10.89		7.46	
Kingman	39.77	49.51	38.35		24.59		33.13		35.56		37.11		41.92	
Palmyra	24.95	31.99	35.07		22.66		25.02		35.35		31.11		42.77	
Wake				31.98		19.29		22.56		31.40		32.34		

Table 8. Mean percent cover of macroalgae from RAMP sites collected from towed-diver surveys in the PRIA

Year	2001	2002	2004	2005	2006	2007	2008	2009	2010	2011	2012	2014	2015	2016
Baker	12.33	2.11	12.63		9.29		8.09		1.60		8.05		2.15	
Howland	2.58	5.34	13.01		3.57		6.14		0.64		6.07		1.08	
Jarvis	28.75	10.88	25.03		38.14		24.01		7.35		7.58		3.94	
Johnston			25.06		6.90		8.82		1.57		8.49		2.49	
Kingman	4.36	5.36	27.04		7.81		7.31		3.97		5.05		2.04	
Palmyra	13.28	10.45	23.14		15.17		11.98		4.76		8.94		4.35	
Wake				22.88		18.74		12.00		8.30		6.80		

Table 9. Mean percent cover of crustose coralline algae from RAMP sites collected from towed-diver surveys in the PRIA

Year	2001	2002	2004	2005	2006	2007	2008	2009	2010	2011	2012	2014	2015	2016
Baker	31.66	37.57	39.61		33.43		23.09		23.40		24.03		32.80	
Howland	36.60	27.40	34.26		22.60		22.59		15.73		18.12		21.25	
Jarvis	29.11	29.56	34.76		24.23		11.82		30.29		24.20		27.48	
Johnston			30.54		19.50		16.07		17.13		17.49		17.45	
Kingman	33.04	16.4	17.49		23.50		13.45		9.20		8.45		9.64	
Palmyra	38.46	24.46	27.26		26.30		18.02		13.87		17.09		10.28	
Wake				1.01		6.43		3.87		4.15		1.13		

2.5.3.3 Oceanography and Water Quality

The water column is also designated as EFH for selected MUS life stages at various depths. For larval stages of all species except deepwater shrimp, the water column is EFH from the shoreline to the EEZ. Coral reef species egg and larval EFH is to a depth of 100 m; crustaceans, 150m; and bottomfish, 400 m. Please see the Climate and Oceanic Indicators section (Section 2.4) for information related to oceanography and water quality.

2.5.4 Report on Review of EFH Information

A review of the biological components of crustacean EFH was originally meant to be included as Appendix C, but will be implemented into the report when available.

2.5.5 EFH Levels

NMFS guidelines codified at 50 C.F.R. § 600.815 recommend Councils organize data used to describe and identify EFH into the following four levels:

- Level 1: Distribution data are available for some or all portions of the geographic range of the species.
- Level 2: Habitat-related densities of the species are available.
- Level 3: Growth, reproduction, or survival rates within habitats are available.
- Level 4: Production rates by habitat are available.

The Council adopted a fifth level, denoted Level 0, for situations in which there is no information available about the geographic extent of a particular managed species' life stage. The existing level of data for individual MUS in each fishery are presented in tables per fishery. Each fishery section also includes the description of EFH, the method used to assess the value of the habitat to the species, description of data sources used if there was analysis, and description of method for analysis. A section summarizing the annual review that was performed follows.

2.5.5.1 Precious Corals

EFH for precious corals was originally designated in Amendment 4 to the Precious Corals Fishery Management Plan (64 FR 19067, 19 April 1999) using the level of data found in Table 10.

Table 10. Level of EFH information available for the Western Pacific precious coral MUS

Species	Pelagic Phase (Larval Stage)	Benthic Phase	Source(s)
Pink Coral (<i>Corallium</i>)			
<i>Pleurocorallium secundum</i> (prev. <i>Corallium secundum</i>)	0	1	Figuroa and Baco (2014); HURL database
<i>C. regale</i>	0	1	HURL database
<i>Hemicorallium laauense</i> (prev. <i>C. laauense</i>)	0	1	HURL database
Gold Coral			
<i>Kulamanamana haumea</i>	0	1	Sinniger et al. (2013); HURL database
<i>Callogorgia gilberti</i>	0	1	HURL database
<i>Narella</i> spp.	0	1	HURL database
Bamboo Coral			
<i>Lepidisis olapa</i>	0	1	HURL database
<i>Acanella</i> spp.	0	1	HURL database
Black Coral			
<i>Antipathes griggsi</i> (prev. <i>Antipathes dichotoma</i>)	0	2	Opresko (2009) HURL database
<i>A. grandis</i>	0	1	HURL database
<i>Myriopathes ulex</i> (prev. <i>A. ulex</i>)	0	1	Opresko (2009) HURL database

2.5.5.2 Bottomfish and Seamount Groundfish

EFH for bottomfish and seamount groundfish was originally designated in Amendment 6 to the Bottomfish and Seamount Groundfish FMP (64 FR 19067, 19 April 1999) using the level of data found in Table 11.

Table 11. Level of EFH information available for the Western Pacific BMUS and seamount groundfish MUS complex

Life History Stage	Eggs	Larvae	Juvenile	Adult
Bottomfish: (scientific/English common)				
<i>Aphareus rutilans</i> (red snapper/silvermouth)	0	0	0	2
<i>Aprion virescens</i> (gray snapper/jobfish)	0	0	1	2
<i>Caranx ignobilis</i> (giant trevally/jack)	0	0	1	2
<i>C. lugubris</i> (black trevally/jack)	0	0	0	2

Life History Stage	Eggs	Larvae	Juvenile	Adult
<i>Epinephelus faciatus</i> (blacktip grouper)	0	0	0	1
<i>E. quernus</i> (sea bass)	0	0	1	2
<i>Etelis carbunculus</i> (red snapper)	0	0	1	2
<i>E. coruscans</i> (red snapper)	0	0	1	2
<i>Lethrinus amboinensis</i> (ambon emperor)	0	0	0	1
<i>L. rubrioperculatus</i> (redgill emperor)	0	0	0	1
<i>Lutjanus kasmira</i> (blueline snapper)	0	0	1	1
<i>Pristipomoides auricilla</i> (yellowtail snapper)	0	0	0	2
<i>P. filamentosus</i> (pink snapper)	0	0	1	2
<i>P. flavipinnis</i> (yelloweye snapper)	0	0	0	2
<i>P. seiboldi</i> (pink snapper)	0	0	1	2
<i>P. zonatus</i> (snapper)	0	0	0	2
<i>Pseudocaranx dentex</i> (thicklip trevally)	0	0	1	2
<i>Seriola dumerili</i> (amberjack)	0	0	0	2
<i>Variola louti</i> (lunartail grouper)	0	0	0	2
Seamount Groundfish:				
<i>Beryx splendens</i> (alfonsin)	0	1	2	2
<i>Hyperoglyphe japonica</i> (ratfish/butterfish)	0	0	0	1
<i>Pseudopentaceros richardsoni</i> (armorhead)	0	1	1	3

2.5.5.3 Crustaceans

EFH for crustaceans MUS was originally designated in Amendment 10 to the Crustaceans FMP (64 FR 19067, 19 April 1999) using the level of data found in Table 12. EFH definitions were also approved for deepwater shrimp through an amendment to the Crustaceans FMP in 2008 (73 FR 70603, 21 November 2008).

Table 12. Level of EFH information available for the Western Pacific CMUS complex

Life History Stage	Eggs	Larvae	Juvenile	Adult
Crustaceans: (english common\scientific)				
Spiny lobster (<i>Panulirus marginatus</i>)	2	1	1-2	2-3
Spiny lobster (<i>Panulirus pencillatus</i>)	1	1	1	2
Common slipper lobster (<i>Scyllarides squammosus</i>)	2	1	1	2-3
Ridgeback slipper lobster (<i>Scyllarides haanii</i>)	2	0	1	2-3
Chinese slipper lobster (<i>Parribacus antarcticus</i>)	2	0	1	2-3
Kona crab (<i>Ranina ranina</i>)	1	0	1	1-2

2.5.5.4 Coral Reef

Essential Fish Habitat for coral reef ecosystem species was originally designated in the Coral Reef Ecosystem FMP (69 FR 8336, 24 February 2004). An EFH review of CREMUS has not

been undertaken, as the Council recently completed its process of re-designating certain CREMUS into the ecosystem component species (ECS) classification. Ecosystem component species do not require EFH designations, as they are not a managed species.

2.5.6 Research and Information Needs

The Council has identified the following scientific data needs to more effectively address the EFH provisions:

2.5.6.1 All FMP Fisheries

- Distribution of early life history stages (eggs and larvae) of management unit species by habitat
- Juvenile habitat (including physical, chemical, and biological features that determine suitable juvenile habitat)
- Food habits (feeding depth, major prey species etc.)
- Habitat-related densities for all MUS life history stages
- Growth, reproduction, and survival rates for MUS within habitats

2.5.6.2 Bottomfish Fishery

- Inventory of marine habitats in the EEZ of the Western Pacific region
- Data to obtain a better SPR estimate for American Samoa's bottomfish complex
- Baseline (virgin stock) parameters (CPUE, percent immature) for the Guam/CNMI deep-water and shallow-water bottomfish complexes
- High resolution maps of bottom topography/currents/water masses/primary productivity
- Habitat utilization patterns for different life history stages and species

2.5.6.3 Crustaceans Fishery

- Identification of post-larval settlement habitat of all CMUS
- Identification of "source/sink" relationships in the NWHI and other regions (i.e. relationships between spawning sites settlement using circulation models, genetic techniques, etc.)
- Establish baseline parameters (e.g., CPUE) for the Guam and Northern Marianas crustacean populations
- Research to determine habitat-related densities for all CMUS life history stages in American Samoa, Guam, Hawaii, and CNMI
- High resolution mapping of bottom topography, bathymetry, currents, substrate types, algal beds, habitat relief

2.5.6.4 Precious Corals Fishery

- Distribution, abundance, and status of precious corals in the PRIAs

2.5.7 References

- 64 FR 19067. Fisheries Off West Coast States and in the Western Pacific; Pelagic Fisheries, Amendment 8; Crustacean Fisheries, Amendment 10; Bottomfish and Seamount Groundfish Fisheries, Amendment 6; Precious Corals Fisheries, Amendment 4, Rule. *Federal Register* 64 (19 April 1999): 19067-19069. Downloaded from <https://www.govinfo.gov/content/pkg/FR-1999-04-19/pdf/99-9728.pdf>.
- 69 FR 8336. Fisheries off West Coast States and in the Western Pacific; Coral Reef Ecosystems Fishery Management Plan for the Western Pacific, Final Rule. *Federal Register* 69 (24 February 2004): 8336-8349. Downloaded from <http://www.wpcouncil.org/precious/Documents/FMP/Amendment5-FR-FinalRule.pdf>.
- 73 FR 70603. Fisheries in the Western Pacific; Crustacean Fisheries; Deepwater Shrimp, Final Rule. *Federal Register* 73 (21 November 2008): 70603-70605. Downloaded from <https://www.govinfo.gov/content/pkg/FR-2008-11-21/pdf/E8-27773.pdf>.
- Brainard R.B., Maragos, J., Schroeder, R., Kenyon, J.C., Vroom PS, Godwin S, Hoeke, R.K., Aeby, G.S., Moffitt, R., Lammers, M., Gove, J., Timmers, M., Holzwarth, S.R., and S. Kolinski, 2005. The state of coral reef ecosystems of the Pacific Remote Island Areas. In: Waddell JE (Ed.). The state of coral reef ecosystems of the United States and Pacific Freely Associated States: 2005, p. 338-372. NOAA Technical Memorandum NOS NCCOS 11. NOAA/NCCOS Center for Coastal Monitoring and Assessment's Biogeography Team. 522 p.
- DesRochers, A., 2016. "Benthic Habitat Mapping." NOAA Fisheries Center, Honolulu, HI. Presentation. 6 April 2016.
- Figueroa, D.F. and A.R. Baco, 2014. Complete mitochondrial genomes elucidate phylogenetic relationships of the deep-sea octocoral families Coralliidae and Paragorgiidae. *Deep Sea Research Part II: Topical Studies in Oceanography*, 99, pp.83-91.
- McCoy, K., Heenan, A., Asher, J., Ayotte, P., Gorospe, K., Gray, A., Lino, K., Zamzowm J., and I. Williams, 2017. Pacific Reef Assessment and Monitoring Program Data Report: Ecological monitoring 2016 - reef fishes and benthic habitats of the main Hawaiian Islands, Northwestern Hawaiian Islands, Pacific Remote Island Areas, and American Samoa. NOAA Pacific Islands Fisheries Science Center. PIFSC Data Report DR-17-001. 66 pp.
- Miller, J., Battista, T., Pritchett, A, Rohmann, S, and J. Rooney, 2011. Coral Reef Conservation Program Mapping Achievements and Unmet Needs. 68 pp.
- Minton, D., 2017. Non-fishing effects that may adversely affect essential fish habitat in the Pacific Islands region, Final Report. NOAA National Marine Fisheries Service, Contract AB-133F-15-CQ-0014. 207 pp.

- Opresko, D.M., 2009. A New Name for the Hawaiian Antipatharian Coral Formerly Known as *Antipathes dichotoma* (Cnidaria: Anthozoa: Antipatharia) 1. *Pacific Science*, 63(2), pp. 277-292.
- PIFSC. 2016. Ecosystem Sciences. Coral Reef Ecosystem Survey Methods. Benthic Monitoring. http://www.pifsc.noaa.gov/cred/survey_methods.php. Updated 1 April 2016.
- PIFSC. 2011. Coral reef ecosystems of American Samoa: a 2002-2010 overview. NOAA Fisheries Pacific Islands Fisheries Science Center, PIFSC Special Publication, SP-11-02, 48 pp.
- Ryan, W.B.F., S.M. Carbotte, S.M., Coplan, J.O., O'Hara, S., Melkonian, A., Arko, R., Weissel, R.A., Ferrini, V., Goodwillie, A., Nitsche, F., Bonczkowski, J., and R. Zemsky, 2009. Global Multi-Resolution Topography synthesis, *Geochem. Geophys. Geosyst.*, 10, Q03014.
- Sinniger, F., Ocana, O.V., and A.R. Baco, 2013. Diversity of zoanthids (Anthozoa: Hexacorallia) on Hawaiian seamounts: description of the Hawaiian gold coral and additional zoanthids. *PloS one*, 8(1), p.e52607.

2.6 MARINE PLANNING

2.6.1 Introduction

Marine planning is a science-based management tool being utilized regionally, nationally and globally to identify and address issues of multiple human uses, ecosystem health and cumulative impacts in the coastal and ocean environment. The Council's efforts to formalize incorporation of marine planning in its actions began in response to Executive Order (EO) 13547, *Stewardship of the Ocean, Our Coasts, and the Great Lakes*. EO 13158, *Marine Protected Areas*, proposes that agencies strengthen the management, protection, and conservation of existing MPAs, develop a national system of MPAs representing diverse ecosystems, and avoid causing harm to MPAs through federal activities. MPAs, or marine managed areas (MMAs) are one tool used in fisheries management and marine planning.

At its 165th meeting in March 2016, in Honolulu, Hawai'i, the Council approved the following objective for the FEPs: Consider the Implications of Spatial Management Arrangements in Council Decision-making. The following sub-objectives apply:

- a. Identify and prioritize research that examines the positive and negative consequences of areas that restrict or prohibit fishing to fisheries, fishery ecosystems, and fishermen, such as the Bottomfish Fishing Restricted Areas (BRFAs), military installations, NWHI restrictions, and Marine Life Conservation Districts (MLCDs).
- b. Establish effective spatially-based fishing zones.
- c. Consider modifying or removing spatial-based fishing restrictions that are no longer necessary or effective in meeting their management objectives.
- d. As needed, periodically evaluate the management effectiveness of existing spatial-based fishing zones in Federal waters.

To monitor implementation of this objective, this annual report includes the Council's spatially-based fishing restrictions or MMAs, the goals associated with those, and the most recent evaluation. Council research needs are not tracked in this report.

To meet the EFH and National Environmental Policy Act (NEPA) mandates, this annual report tracks activities that occur in the ocean that are of interest to the Council, and incidents or facilities that may contribute to cumulative impact. NMFS is responsible for NEPA compliance, and the Council must assess the environmental effects of ocean activities for the EFH cumulative impacts section of the FEP.

2.6.1.1 Response to Previous Council Recommendations

There are no standing Council recommendations indicating review deadlines for PRIA MMAs.

2.6.1.2 MMAs established under FMPs

Council-established MMAs were compiled from 50 CFR § 665, Western Pacific Fisheries, the Federal Register, and Council amendment documents. All regulated fishing areas and large MMAs, including the Pacific Remote Islands Marine National Monument (PRIMNM), are shown in Figure 35.




Figure 35. Regulated fishing areas of the PRIAs

Table 13. MMAs established under FEPs from [50 CFR § 665](#)

Name	FEP	Island	50 CFR /FR /Amendment Reference	Marine Area (km ²)	Fishing Restriction	Goals	Most Recent Evaluation	Review Deadline
Howland Island No-Take MPA/PRI Marine National Monument	PRIA/Pelagic	Howland Island	665.599 and 665.799(a)(1) 69 FR 8336 Coral Reef Ecosystem FEP 78 FR 32996 PRIA FEP Am. 2	-	All Take Prohibited	Minimize adverse human impacts on coral reef resources; commercial fishing prohibited within 12 nm.	2013	-
Jarvis Island No-Take MPA/PRI Marine National Monument	PRIA/Pelagic	Jarvis Island	665.599 and 665.799(a)(1) 69 FR 8336 Coral Reef Ecosystem FEP 78 FR 32996 PRIA FEP Am. 2	-	All Take Prohibited	Minimize adverse human impacts on coral reef resources; commercial fishing prohibited within 12 nm.	2013	-
Baker Island No-Take MPA/PRI Marine National Monument	PRIA/Pelagic	Baker Island	665.599 and 665.799(a)(1) 69 FR 8336 Coral Reef Ecosystem FEP 78 FR 32996 PRIA FEP Am. 2	-	All Take Prohibited	Minimize adverse human impacts on coral reef resources; commercial fishing prohibited within 12 nm.	2013	-

Name	FEP	Island	50 CFR /FR /Amendment Reference	Marine Area (km ²)	Fishing Restriction	Goals	Most Recent Evaluation	Review Deadline
Kingman Reef No-Take MPA/PRI Marine National Monument	PRIA/Pelagic	Kingman Reef	665.599 and 665.799(a)(1) 69 FR 8336 Coral Reef Ecosystem FEP 78 FR 32996 PRIA FEP Am. 2	-	All Take Prohibited	Minimize adverse human impacts on coral reef resources; all fishing prohibited within 12 nm.	2013	-
Johnston Atoll Low-Use MPA/PRI Marine National Monument	PRIA/Pelagic	Johnston Atoll	69 FR 8336 Coral Reef Ecosystem FEP 78 FR 32996 PRIA FEP Am. 2	-	Special Permit Only	Minimize adverse human impacts on coral reef resources; superseded by prohibiting fishing within 12 nm in Am. 2.	2013	-
Palmyra Atoll Low-Use MPAs/PRI Marine National Monument	PRIA/Pelagic	Palmyra Atoll	69 FR 8336 Coral Reef Ecosystem FEP 78 FR 32996 PRIA FEP Am. 2	-	Special Permit Only	Minimize adverse human impacts on coral reef resources; superseded by prohibiting fishing within 12 nm in Am. 2.	2013	-
Wake Island Low-Use MPA/PRI Marine National Monument	PRIA/Pelagic	Wake Island	69 FR 8336 Coral Reef Ecosystem FEP 78 FR 32996 PRIA FEP Am. 2	-	Special Permit Only	Minimize adverse human impacts on coral reef resources; superseded by prohibiting fishing within 12 nm in Am. 2.	2013	-

2.6.2 Activities and Facilities

There are no aquaculture, alternative energy facilities, or military training and testing activities occurring in the US EEZ around the PRIAs at this time. The Plan Team will add to this section as new facilities are proposed and/or built.

2.6.3 Pacific Islands Regional Planning Body Report

In June 2018, President Trump signed the EO 13840 *Regarding the Ocean Policy to Advance Economic, Security, and Environmental Interests of the United States*, which revoked EO 13547. The new EO eliminated the mandate for the federal government to participate in ocean planning at a regional level and eliminated the regional planning bodies. As such, the Pacific Islands Regional Planning Body (RPB) no longer exists and ocean planning will now occur at a local level led by Hawaii and the territories.

However, EO 13840 established a policy focused on public access to marine data and information, and requires federal agencies to 1) coordinate activities regarding ocean-related matters and 2) facilitate the coordination and collaboration of ocean-related matters with governments and ocean stakeholders. To that end, the [American Samoa Coastal and Marine Spatial Planning Data Portal](#) was created by Marine Cadastre (<https://marinecadastre.gov/viewers/>). The intent is for it to be expanded to include the Marianas, PRIAs, and Hawaii, and be titled the Pacific Islands Regional Marine Planner.

2.6.4 References

- 50 CFR § 665. Fisheries in the Western Pacific. Title 50 *Code of Federal Regulations*, Pt. 665. Electronic Code of Federal Regulations data current as of March 16, 2016. Viewed at http://www.ecfr.gov/cgi-bin/retrieveECFR?gp=&SID=b28abb7da3229173411daf43959fcbd1&n=50y13.0.1.1.2&r=PART&ty=HTML#_top.
- 56 FR 52214. Pelagic Fisheries of the Western Pacific Region, Final Rule. *Federal Register* 56 (18 October 1991): 52214-52217. Downloaded from <http://www.wpcouncil.org/pelagic/Documents/FMP/Amendment3-FR-FinalRule.pdf>.
- 57 FR 7661. Pelagic Fisheries of the Western Pacific Region, Final Rule. *Federal Register* 57 (4 March 1992): 7661-7665. Downloaded from <http://www.wpcouncil.org/pelagic/Documents/FMP/Amendment5-FR-FinalRule.pdf>.
- 69 FR 8336. Fisheries off West Coast States and in the Western Pacific; Coral Reef Ecosystems Fishery Management Plan for the Western Pacific, Final Rule. *Federal Register* 69 (24 February 2004): 8336-8349. Downloaded from <http://www.wpcouncil.org/precious/Documents/FMP/Amendment5-FR-FinalRule.pdf>.
- 78 FR 32996. Western Pacific Fisheries; Fishing in the Marianas Trench, Pacific Remote Islands, and Rose Atoll Marine National Monuments, Final Rule. *Federal Register* 78 (3 June 2013): 32996-33007. Downloaded from <http://www.wpcouncil.org/precious/Documents/FMP/Amendment5-FR-FinalRule.pdf>.

3 DATA INTEGRATION

The purpose of this section (“Chapter 3”) of the Annual SAFE Report is to identify and evaluate potential fishery ecosystem relationships between fishery parameters and ecosystem variables to assess how changes in the ecosystem can affect fisheries across the Western Pacific region. “Fishery ecosystem relationships” are those associations between various fishery-dependent data measures (e.g., catch, effort, or CPUE), and other environmental attributes (e.g., temperature, precipitation, current velocity) that may contribute to observed trends or act as potential indicators of the status of prominent stocks in the fishery. Data integration analyses represent a first step in a sequence of exploratory analyses that will be utilized to inform new assessments of what factors may be useful going forward, and were first incorporated in the 2017 versions of the Annual SAFE Reports.

To support the development of Chapter 3 of the annual SAFE report, staff from the Council, NMFS PIFSC and PIRO, and Triton Aquatics (consultants), held a SAFE Report Data Integration Workshop (hereafter, “the Workshop”) on November 30, 2016 to identify potential fishery ecosystem relationships relevant to local policy in the Western Pacific region and determine appropriate methods to analyze them. The archipelagic fisheries group developed nearly 30 potential fishery ecosystem relationships to examine across bottomfish, coral reef, and crustacean fisheries based on data reliability, suitability of methodology, repeatability on an annual basis, and how well analyses could potentially inform management decisions (Table 14). It is important to note that these lists were developed before the ecosystem component FEP amendment was developed.

Table 14. List of brainstormed potential archipelagic island fishery relationships scored and ranked from highest to lowest priority

Relationships	FEP	Score	Rank
Bottomfish catch/effort/CPUE/species composition and benthos/substrate (i.e., depth, structure)	All	22	3
Bottomfish catch/effort/ CPUE /species composition and Pacific Decadal Oscillation	All	20	3
Coral reef fish/fishery/biomass and temperature-derived variable	All	20	3
Akule/opelu and precipitation (MHI and Guam)	HI	20	3
Bottomfish catchability and wind speed	All	19	3
Coral reef fish/fishery/biomass and chlorophyll- <i>a</i> (with phase lag)	All	19	3
Bottomfish Catch /CPUE and lunar cycle/moon phase	All	19	3
Bottomfish catch/effort/ CPUE /species composition and sea-level height (eddy feature)	All	18	2
Coral reef fish/fishery/biomass and Pacific Decadal Oscillation	All	18	2
Green/red spiny lobster catch/CPUE and vertical relief	HI	18	2

Green/red spiny lobster catch/CPUE and Pacific Decadal Oscillation	HI	18	2
Bottomfish catchability and fishing conditions (i.e. surface, subsurface current, speed, and direction)	All	17	2
Coral reef fish/fishery/biomass and moon phase	All	17	2
Coral reef fish/fishery/biomass and Oceanic Niño Index	All	17	2
Coral reef fish/fishery/biomass and sea-level height	All	17	2
Coral reef fish/fishery/biomass and pH	All	17	2
Bottomfish catch/effort/ CPUE /species composition and temperature-derived variable (e.g. temperature at depth)	All	16	2
Bottomfish catch/effort/ CPUE /species composition and chlorophyll- <i>a</i> (with phase lag)	All	16	2
Bottomfish catch/effort/ CPUE /species composition and precipitation	All	16	2
Coral reef fish/fishery/biomass and structural complexity /benthic habitat	All	16	2
Bottomfish catch/effort/ CPUE /species composition and dissolved oxygen	All	15	2
Coral reef fish/fishery/biomass and precipitation	All	14	2
Bottomfish catch/effort/ CPUE /species composition and pH	All	13	2
Bottomfish catch/effort/ CPUE /species composition and predator abundance	All	12	2
Coral reef fish/fishery/biomass and salinity	All	12	2
Coral reef fish/fishery/biomass and dissolved oxygen	All	12	2
Bottomfish catch/effort/ CPUE /species composition and salinity	All	10	1

The data integration chapter of this report is still under development. The archipelagic data integration chapter explores the potential association between fishery parameters and ecologically-associated variables that may be able to explain a portion of the variance in fishery-dependent data. The Workshop produced a long list of fishery and ecosystem variable combinations that comprise a significant workload that the participants could not take on without sufficient data coverage. Though a contractor completed exploratory evaluations for the MHI, Guam, CNMI, and American Samoa in 2017 for inclusion in the 2017 Annual SAFE Report, no explicit analyses were conducted for the PRIAs. Going forward with the data integration analyses and presentation of results for Chapter 3 of the Annual SAFE Reports, the Plan Team suggested several improvements to implement in the coming year: standardizing and correcting values in CPUE time series, incorporating longer stretches of phase lag, completing comparisons on the species-level and by dominant gear types, incorporating local knowledge on shifts in fishing dynamics over the course of the time series, and utilizing the exact environmental data sets presented in the ecosystem consideration chapter of the annual report. However, no data integration analyses could be completed for the PRIAs in the 2018 Annual SAFE Report. Many

of these recommendations were applied to datasets from Hawaii in 2018, and will similarly be done for PRIA data integration analyses in the upcoming report cycles if the fisheries there become more active and more fishery-dependent data becomes available. Implementation of these suggestions will allow for the preparation and publication of a more finalized version of the data integration chapter in coming years as data availability allows.

APPENDIX A: LIST OF MANAGEMENT UNIT SPECIES

PRIA

The PRIA species list and FSSI status will be made available in subsequent reports as resources allow. Please see the PRIA FEP and implementing regulations for the list of managed species.

APPENDIX B: LIST OF PROTECTED SPECIES AND DESIGNATED CRITICAL HABITAT

Table B-1. Protected species found or reasonably believed to be found near or in Hawai'i shallow-set longline waters.

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
Seabirds					
Laysan Albatross	<i>Phoebastria immutabilis</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Black-Footed Albatross	<i>Phoebastria nigripes</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Short-Tailed Albatross	<i>Phoebastria albatrus</i>	Endangered	N/A	Breeding visitor in the NWHI	35 FR 8495, 65 FR 46643, Pyle & Pyle 2009
Northern Fulmar	<i>Fulmarus glacialis</i>	Not Listed	N/A	Winter resident	Pyle & Pyle 2009
Kermadec Petrel	<i>Pterodroma neglecta</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Herald Petrel	<i>Pterodroma arminjoniana</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Murphy's Petrel	<i>Pterodroma ultima</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Mottled Petrel	<i>Pterodroma inexpectata</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Juan Fernandez Petrel	<i>Pterodroma externa</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Hawaiian Petrel	<i>Pterodroma sandwichensis</i> (<i>Pterodroma phaeopygia sandwichensis</i>)	Endangered	N/A	Breeding visitor in the MHI	32 FR 4001, Pyle & Pyle 2009
White-Necked Petrel	<i>Pterodroma cervicalis</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Bonin Petrel	<i>Pterodroma hypoleuca</i>	Not Listed	N/A	Breeding visitor in the NWHI	Pyle & Pyle 2009
Black-Winged Petrel	<i>Pterodroma nigripennis</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Cook Petrel	<i>Pterodroma cookii</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Stejneger Petrel	<i>Pterodroma longirostris</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Pycroft Petrel	<i>Pterodroma pycrofti</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Bulwer Petrel	<i>Bulweria bulwerii</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Flesh-Footed Shearwater	<i>Ardenna carneipes</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Wedge-Tailed Shearwater	<i>Ardenna pacifica</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Buller's Shearwater	<i>Ardenna bulleri</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
Sooty Shearwater	<i>Ardenna grisea</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Short-Tailed Shearwater	<i>Ardenna tenuirostris</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Christmas Shearwater	<i>Puffinus nativitatis</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Newell's Shearwater	<i>Puffinus newelli</i> (<i>Puffinus auricularis newelli</i>)	Threatened	N/A	Breeding visitor	40 FR 44149, Pyle & Pyle 2009
Wilson's Storm-Petrel	<i>Oceanites oceanicus</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Leach's Storm-Petrel	<i>Oceanodroma leucorhoa</i>	Not Listed	N/A	Winter resident	Pyle & Pyle 2009
Band-Rumped Storm-Petrel	<i>Oceanodroma castro</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Tristram Storm-Petrel	<i>Oceanodroma tristrami</i>	Not Listed	N/A	Breeding visitor in the NWHI	Pyle & Pyle 2009
White-Tailed Tropicbird	<i>Phaethon lepturus</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Red-Tailed Tropicbird	<i>Phaethon rubricauda</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Masked Booby	<i>Sula dactylatra</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Brown Booby	<i>Sula leucogaster</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Red-Footed Booby	<i>Sula sula</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Great Frigatebird	<i>Fregata minor</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Lesser Frigatebird	<i>Fregata ariel</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Laughing Gull	<i>Leucophaeus atricilla</i>	Not Listed	N/A	Winter resident in the MHI	Pyle & Pyle 2009
Franklin Gull	<i>Leucophaeus pipixcan</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Ring-Billed Gull	<i>Larus delawarensis</i>	Not Listed	N/A	Winter resident in the MHI	Pyle & Pyle 2009
Herring Gull	<i>Larus argentatus</i>	Not Listed	N/A	Winter resident in the NWHI	Pyle & Pyle 2009
Slaty-Backed Gull	<i>Larus schistisagus</i>	Not Listed	N/A	Winter resident in the NWHI	Pyle & Pyle 2009
Glaucous-Winged Gull	<i>Larus glaucescens</i>	Not Listed	N/A	Winter resident	Pyle & Pyle 2009
Brown Noddy	<i>Anous stolidus</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Black Noddy	<i>Anous minutus</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Blue-Gray Noddy	<i>Procelsterna cerulea</i>	Not Listed	N/A	Breeding visitor in the NWHI	Pyle & Pyle 2009
White Tern	<i>Gygis alba</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
Sooty Tern	<i>Onychoprion fuscatus</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Gray-Backed Tern	<i>Onychoprion lunatus</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Little Tern	<i>Sternula albifrons</i>	Not Listed	N/A	Breeding visitor in the NWHI	Pyle & Pyle 2009
Least Tern	<i>Sternula antillarum</i>	Not Listed	N/A	Breeding visitor in the NWHI	Pyle & Pyle 2009
Arctic Tern	<i>Sterna paradisaea</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
South Polar Skua	<i>Stercorarius maccormicki</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Pomarine Jaeger	<i>Stercorarius pomarinus</i>	Not Listed	N/A	Winter resident in the MHI	Pyle & Pyle 2009
Parasitic Jaeger	<i>Stercorarius parasiticus</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Long-Tailed Jaeger	<i>Stercorarius longicaudus</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Sea turtles					
Green Sea Turtle	<i>Chelonia mydas</i>	Threatened (Central North Pacific DPS)	N/A	Most common turtle in the Hawaiian Islands, much more common in nearshore state waters (foraging grounds) than offshore federal waters. Most nesting occurs on French Frigate Shoals in the NWHI. Foraging and haul out in the MHI.	43 FR 32800, 81 FR 20057, Balazs et al. 1992, Kolinski et al. 2001
Green Sea Turtle	<i>Chelonia mydas</i>	Threatened (East Pacific DPS)	N/A	Nest primarily in Mexico and the Galapagos Islands. Little known about their pelagic range west of 90°W, but may range as far as the Marshall Islands. Genetic testing confirmed that they are incidentally taken in the HI DSLL fishery.	43 FR 32800, 81 FR 20057, WPRFMC 2009, Clifton et al. 1982, Karl & Bowen 1999
Hawksbill Sea Turtle	<i>Eretmochelys imbricata</i>	Endangered ^a	N/A	Small population foraging around Hawai'i and low level nesting on Maui and Hawai'i Islands. Occur worldwide in tropical and subtropical waters.	35 FR 8491, NMFS & USFWS 2007, Balazs et al. 1992, Katahira et al. 1994
Leatherback Sea Turtle	<i>Dermochelys coriacea</i>	Endangered ^a	N/A	Regularly sighted in offshore waters, especially at the southeastern end of the archipelago.	35 FR 8491, NMFS & USFWS 1997

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
Loggerhead Sea Turtle	<i>Caretta caretta</i>	Endangered (North Pacific DPS)	N/A	Rare in Hawai'i. Found worldwide along continental shelves, bays, estuaries and lagoons of tropical, subtropical, and temperate waters.	43 FR 32800, 76 FR 58868, Dodd 1990, Balazs 1979
Olive Ridley Sea Turtle	<i>Lepidochelys olivacea</i>	Threatened (Entire species, except for the breeding population on the Pacific coast of Mexico, which is listed as endangered)	N/A	Rare in Hawai'i. Occurs worldwide in tropical and warm temperate ocean waters.	43 FR 32800, Pitman 1990, Balacz 1982
Marine mammals					
Blainville's Beaked Whale	<i>Mesoplodon densirostris</i>	Not Listed	Non-strategic	Found worldwide in tropical and temperate waters	Mead 1989
Blue Whale	<i>Balaenoptera musculus</i>	Endangered	Strategic	Acoustically recorded off of Oahu and Midway Atoll, small number of sightings around Hawai'i. Considered extremely rare, generally occur in winter and summer.	35 FR 18319, Bradford et al. 2013, Northrop et al. 1971, Thompson & Friedl 1982, Stafford et al. 2001
Bottlenose Dolphin	<i>Tursiops truncatus</i>	Not Listed	Non-strategic	Distributed worldwide in tropical and warm-temperate waters. Pelagic stock distinct from island-associated stocks.	Perrin et al. 2009, Martien et al. 2012
Bryde's Whale	<i>Balaenoptera edeni</i>	Not Listed	Unknown	Distributed widely across tropical and warm-temperate Pacific Ocean.	Leatherwood et al. 1982
Common Dolphin	<i>Delphinus delphis</i>	Not Listed	N/A	Found worldwide in temperate and subtropical seas.	Perrin et al. 2009
Cuvier's Beaked Whale	<i>Ziphius cavirostris</i>	Not Listed	Non-strategic	Occur year round in Hawaiian waters.	McSweeney et al. 2007
Dall's Porpoise	<i>Phocoenoides dalli</i>	Not Listed	Non-strategic	Range across the entire north Pacific Ocean.	Hall 1979
Dwarf Sperm Whale	<i>Kogia sima</i>	Not Listed	Non-strategic	Most common in waters between 500 m and 1,000 m in depth. Found worldwide in tropical and warm-temperate waters.	Nagorsen 1985, Baird et al. 2013

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
False Killer Whale	<i>Pseudorca crassidens</i>	Not Listed	Non-strategic	Found worldwide in tropical and warm-temperate waters. Pelagic stock tracked to within 11 km of Hawaiian islands.	Stacey et al. 1994, Baird et al. 2012, Bradford et al. 2015
Fin Whale	<i>Balaenoptera physalus</i>	Endangered	Strategic	Infrequent sightings in Hawai'i waters. Considered rare in Hawai'i, though may migrate into Hawaiian waters during fall/winter based on acoustic recordings.	35 FR 18319, Hamilton et al. 2009, Thompson & Friedl 1982
Fraser's Dolphin	<i>Lagenodelphis hosei</i>	Not Listed	Non-strategic	Found worldwide in tropical waters.	Perrin et al. 2009
Guadalupe Fur Seal	<i>Arctocephalus townsendi</i>	Threatened	Strategic	Extremely rare sightings. Little known about their pelagic distribution. Breed mainly on Isla Guadalupe, Mexico.	50 FR 51252, Gallo-Reynoso et al. 2008, Fleischer 1987
Hawaiian Monk Seal	<i>Neomonachus schauinslandi</i>	Endangered ^a	Strategic	Endemic tropical seal. Occurs throughout the archipelago. MHI population spends some time foraging in federal waters during the day.	41 FR 51611, Baker et al. 2011
Humpback Whale	<i>Megaptera novaeangliae</i>	Delisted Due to Recovery (Hawai'i DPS)	Strategic	Migrate through the archipelago and breed during the winter. Common during winter months, when they are generally found within the 100 m isobath.	35 FR 18319, 81 FR 62259, Childerhouse et al. 2008, Wolman & Jurasz 1976, Herman & Antinaja 1977, Rice & Wolman 1978
Killer Whale	<i>Orcinus orca</i>	Not Listed	Non-strategic	Rare in Hawai'i. Prefer colder waters within 800 km of continents.	Mitchell 1975, Baird et al. 2006
Longman's Beaked Whale	<i>Indopacetus pacificus</i>	Not Listed	Non-strategic	Found in tropical waters from the eastern Pacific westward through the Indian Ocean to the eastern coast of Africa. Rare in Hawai'i.	Dalebout 2003, Baird et al. 2013
Melon-Headed Whale	<i>Peponocephala electra</i>	Not Listed	Non-strategic	Found in tropical and warm-temperate waters worldwide, found primarily in equatorial waters. Uncommon in Hawai'i.	Perryman et al. 1994, Barlow 2006, Bradford et al. 2013

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
Minke Whale	<i>Balaenoptera acutorostrata</i>	Not Listed	Non-strategic	Occur seasonally around Hawai i	Barlow 2003, Rankin & Barlow 2005
North Pacific Right Whale	<i>Eubalaena japonica</i>	Endangered ^a	Strategic	Extremely rare in Hawai i waters	35 FR 18319, 73 FR 12024, Rowntree et al. 1980, Herman et al. 1980
Northern Elephant Seal	<i>Mirounga angustirostris</i>	Not Listed	Non-strategic	Females migrate to central North Pacific to feed on pelagic prey.	Le Beouf et al. 2000
Northern Fur Seal	<i>Callorhinus ursinus</i>	Not Listed	Non-strategic	Occur throughout the North Pacific Ocean.	Gelatt et al. 2015
Pacific White-Sided Dolphin	<i>Lagenorhynchus obliquidens</i>	Not Listed	Non-strategic	Endemic to temperate waters of North Pacific Ocean. Occur both on the high seas and along continental margins.	Brownell et al. 1999
Pantropical Spotted Dolphin	<i>Stenella attenuata attenuata</i>	Not Listed	Non-strategic	Common and abundant throughout the Hawaiian archipelago. Pelagic stock occurs outside of insular stock areas (20 km for Oahu and 4-island stocks, 65 km for Hawai i Island stock).	Baird et al. 2013, Oleson et al. 2013
Pygmy Killer Whale	<i>Feresa attenuata</i>	Not Listed	Non-strategic	Small resident population in Hawaiian waters. Found worldwide in tropical and subtropical waters.	McSweeney et al. 2009, Ross & Leatherwood 1994
Pygmy Sperm Whale	<i>Kogia breviceps</i>	Not Listed	Non-strategic	Found worldwide in tropical and warm-temperate waters.	Caldwell & Caldwell 1989
Risso's Dolphin	<i>Grampus griseus</i>	Not Listed	Non-strategic	Found in tropical to warm-temperate waters worldwide.	Perrin et al. 2009
Rough-Toothed Dolphin	<i>Steno bredanensis</i>	Not Listed	Non-strategic	Found in tropical to warm-temperate waters worldwide. Occasionally found offshore of Hawai i.	Perrin et al. 2009, Baird et al. 2013, Barlow 2006, Bradford et al. 2013
Sei Whale	<i>Balaenoptera borealis</i>	Endangered	Strategic	Rare in Hawai i. Generally found in offshore temperate waters.	35 FR 18319, Barlow 2003, Bradford et al. 2013

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
Short-Finned Pilot Whale	<i>Globicephala macrorhynchus</i>	Not Listed	Non-strategic	Found in tropical to warm-temperate waters worldwide. Commonly observed around MHI and present around NWHI.	Shallenberger 1981, Baird et al. 2013, Bradford et al. 2013
Sperm Whale	<i>Physeter macrocephalus</i>	Endangered	Strategic	Found in tropical to polar waters worldwide, most abundant cetaceans in the region. Sighted off the NWHI and the MHI.	35 FR 18319, Rice 1960, Lee 1993, Barlow 2006, Mobley et al. 2000, Shallenberger 1981
Spinner Dolphin	<i>Stenella longirostris</i>	Not Listed	Non-strategic	Found worldwide in tropical and warm-temperate waters. Pelagic stock found outside of island-associated boundaries (10 nm).	Perrin et al. 2009
Striped Dolphin	<i>Stenella coeruleoalba</i>	Not Listed	Non-strategic	Found in tropical to warm-temperate waters throughout the world.	Perrin et al. 2009
Elasmobranchs					
Giant manta ray	<i>Manta birostris</i>	Threatened	N/A	Found worldwide in tropical, subtropical, and temperate waters. Commonly found in upwelling zones, oceanic island groups, offshore pinnacles and seamounts, and on shallow reefs.	Dewar et al. 2008, Marshall et al. 2009, Marshall et al. 2011.
Oceanic whitetip shark	<i>Carcharhinus longimanus</i>	Threatened	N/A	Found worldwide in open ocean waters from the surface to 152 m depth. It is most commonly found in waters > 20°C	Bonfil et al. 2008, Backus et al, 1956, Strasburg 1958, Compagno 1984
Scalloped hammerhead shark	<i>Sphyrna lewini</i>	Endangered (Eastern Pacific DPS)	N/A	Found in coastal areas from southern California to Peru.	Compagno 1984, Baum et al. 2007, Bester 2011
Scalloped hammerhead	<i>Sphyrna lewini</i>	Threatened (Indo-West Pacific DPS)	N/A	Occur over continental and insular shelves, and adjacent deep waters, but rarely found in waters < 22°C. Range from the intertidal and surface to depths up to 450–512 m.	Compagno 1984, Schulze-Haugen & Kohler 2003, Sanches 1991, Klimley 1993
Corals					

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
N/A	<i>Acropora globiceps</i>	Threatened	N/A	Not confirmed in Hawai i waters. Occur on upper reef slopes, reef flats, and adjacent habitats in depths ranging from 0 to 8 m	Veron 2014
N/A	<i>Acropora jacquelineae</i>	Threatened	N/A	Not confirmed in Hawai i waters. Found in numerous subtidal reef slope and back-reef habitats, including but not limited to, lower reef slopes, walls and ledges, mid-slopes, and upper reef slopes protected from wave action, and depth range is 10 to 35 m.	Veron 2014
N/A	<i>Acropora retusa</i>	Threatened	N/A	Not confirmed in Hawai i waters. Occur in shallow reef slope and back-reef areas, such as upper reef slopes, reef flats, and shallow lagoons, and depth range is 1 to 5 m.	Veron 2014
N/A	<i>Acropora speciosa</i>	Threatened	N/A	Not confirmed in Hawai i waters. Found in protected environments with clear water and high diversity of <i>Acropora</i> and steep slopes or deep, shaded waters. Depth range is 12 to 40 meters, and have been found in mesophotic habitat (40-150 m).	Veron 2014
N/A	<i>Euphyllia paradivisa</i>	Threatened	N/A	Not confirmed in Hawai i waters. Found in environments protected from wave action on at least upper reef slopes, mid-slope terraces, and lagoons in depths ranging from 2 to 25 m depth.	Veron 2014
N/A	<i>Isopora crateriformis</i>	Threatened	N/A	Not confirmed in Hawai i waters. Found in shallow, high-wave energy environments, from low tide to at least 12 meters deep, and have been reported from mesophotic depths (less than 50 m depth).	Veron 2014

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
N/A	<i>Seriatopora aculeata</i>	Threatened	N/A	Not confirmed in Hawai'i waters. Found in broad range of habitats including, but not limited to, upper reef slopes, mid-slope terraces, lower reef slopes, reef flats, and lagoons, and depth ranges from 3 to 40 m.	Veron 2014
Invertebrates					
Chambered nautilus	<i>Nautilus pompilius</i>	Threatened	N/A	Found in small, isolated populations throughout the Indo-Pacific on steep-sloped forereefs with sandy, silty, or muddy bottom substrates from depths of 100 m to 500 m.	83 FR 48948, CITES 2016

^a These species have critical habitat designated under the ESA. See Table B-4.

Table B-2. Protected species found or reasonably believed to be found near or in Hawai'i deep-set longline waters.

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
Seabirds					
Laysan Albatross	<i>Phoebastria immutabilis</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Black-Footed Albatross	<i>Phoebastria nigripes</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Short-Tailed Albatross	<i>Phoebastria albatrus</i>	Endangered	N/A	Breeding visitor in the NWHI	35 FR 8495, 65 FR 46643, Pyle & Pyle 2009
Northern Fulmar	<i>Fulmarus glacialis</i>	Not Listed	N/A	Winter resident	Pyle & Pyle 2009
Kermadec Petrel	<i>Pterodroma neglecta</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Herald Petrel	<i>Pterodroma arminjoniana</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Murphy's Petrel	<i>Pterodroma ultima</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Mottled Petrel	<i>Pterodroma inexpectata</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Juan Fernandez Petrel	<i>Pterodroma externa</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Hawaiian Petrel	<i>Pterodroma sandwichensis</i> (<i>Pterodroma phaeopygia sandwichensis</i>)	Endangered	N/A	Breeding visitor in the MHI	32 FR 4001, Pyle & Pyle 2009
White-Necked Petrel	<i>Pterodroma cervicalis</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Bonin Petrel	<i>Pterodroma hypoleuca</i>	Not Listed	N/A	Breeding visitor in the NWHI	Pyle & Pyle 2009
Black-Winged Petrel	<i>Pterodroma nigripennis</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Cook Petrel	<i>Pterodroma cookii</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Stejneger Petrel	<i>Pterodroma longirostris</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Pycroft Petrel	<i>Pterodroma pycrofti</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Bulwer Petrel	<i>Bulweria bulwerii</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Wedge-Tailed Shearwater	<i>Ardenna pacifica</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Buller's Shearwater	<i>Ardenna bulleri</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Sooty Shearwater	<i>Ardenna grisea</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Short-Tailed Shearwater	<i>Ardenna tenuirostris</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Christmas Shearwater	<i>Puffinus nativitatis</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
Newell's Shearwater	<i>Puffinus newelli</i> (<i>Puffinus auricularis newelli</i>)	Threatened	N/A	Breeding visitor	40 FR 44149, Pyle & Pyle 2009
Wilson's Storm-Petrel	<i>Oceanites oceanicus</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Leach's Storm-Petrel	<i>Oceanodroma leucorhoa</i>	Not Listed	N/A	Winter resident	Pyle & Pyle 2009
Band-Rumped Storm-Petrel	<i>Oceanodroma castro</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Tristram Storm-Petrel	<i>Oceanodroma tristrami</i>	Not Listed	N/A	Breeding visitor in the NWHI	Pyle & Pyle 2009
White-Tailed Tropicbird	<i>Phaethon lepturus</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Red-Tailed Tropicbird	<i>Phaethon rubricauda</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Masked Booby	<i>Sula dactylatra</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Nazca Booby	<i>Sula granti</i>	Not Listed	N/A	Vagrant	Pyle & Pyle 2009
Brown Booby	<i>Sula leucogaster</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Red-Footed Booby	<i>Sula sula</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Great Frigatebird	<i>Fregata minor</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Lesser Frigatebird	<i>Fregata ariel</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Laughing Gull	<i>Leucophaeus atricilla</i>	Not Listed	N/A	Winter resident in the MHI	Pyle & Pyle 2009
Franklin Gull	<i>Leucophaeus pipixcan</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Ring-Billed Gull	<i>Larus delawarensis</i>	Not Listed	N/A	Winter resident in the MHI	Pyle & Pyle 2009
Herring Gull	<i>Larus argentatus</i>	Not Listed	N/A	Winter resident in the NWHI	Pyle & Pyle 2009
Slaty-Backed Gull	<i>Larus schistisagus</i>	Not Listed	N/A	Winter resident in the NWHI	Pyle & Pyle 2009
Glaucous-Winged Gull	<i>Larus glaucescens</i>	Not Listed	N/A	Winter resident	Pyle & Pyle 2009
Brown Noddy	<i>Anous stolidus</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Black Noddy	<i>Anous minutus</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Blue-Gray Noddy	<i>Procelsterna cerulea</i>	Not Listed	N/A	Breeding visitor in the NWHI	Pyle & Pyle 2009
White Tern	<i>Gygis alba</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Sooty Tern	<i>Onychoprion fuscatus</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009
Gray-Backed Tern	<i>Onychoprion lunatus</i>	Not Listed	N/A	Breeding visitor	Pyle & Pyle 2009

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
Little Tern	<i>Sternula albifrons</i>	Not Listed	N/A	Breeding visitor in the NWHI	Pyle & Pyle 2009
Least Tern	<i>Sternula antillarum</i>	Not Listed	N/A	Breeding visitor in the NWHI	Pyle & Pyle 2009
Arctic Tern	<i>Sterna paradisaea</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
South Polar Skua	<i>Stercorarius maccormicki</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Pomarine Jaeger	<i>Stercorarius pomarinus</i>	Not Listed	N/A	Winter resident in the MHI	Pyle & Pyle 2009
Parasitic Jaeger	<i>Stercorarius parasiticus</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Long-Tailed Jaeger	<i>Stercorarius longicaudus</i>	Not Listed	N/A	Migrant	Pyle & Pyle 2009
Sea turtles					
Green Sea Turtle	<i>Chelonia mydas</i>	Threatened (Central North Pacific DPS)	N/A	Most common turtle in the Hawaiian Islands, much more common in nearshore state waters (foraging grounds) than offshore federal waters. Most nesting occurs on French Frigate Shoals in the NWHI. Foraging and haulout in the MHI.	43 FR 32800, 81 FR 20057, Balazs et al. 1992, Kolinski et al. 2001
Green Sea Turtle	<i>Chelonia mydas</i>	Threatened (East Pacific DPS)	N/A	Nest primarily in Mexico and the Galapagos Islands. Little known about their pelagic range west of 90°W, but may range as far as the Marshall Islands. Genetic testing confirmed that they are incidentally taken in the HI DSLL fishery.	43 FR 32800, 81 FR 20057, WPRFMC 2009, Clifton et al. 1982, Karl & Bowen 1999
Hawksbill Sea Turtle	<i>Eretmochelys imbricata</i>	Endangered ^a	N/A	Small population foraging around Hawai'i and low level nesting on Maui and Hawai'i Islands. Occur worldwide in tropical and subtropical waters.	35 FR 8491, NMFS & USFWS 2007, Balazs et al. 1992, Katahira et al. 1994
Leatherback Sea Turtle	<i>Dermochelys coriacea</i>	Endangered ^a	N/A	Regularly sighted in offshore waters, especially at the southeastern end of the archipelago.	35 FR 8491, NMFS & USFWS 1997

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
Loggerhead Sea Turtle	<i>Caretta caretta</i>	Endangered (North Pacific DPS)	N/A	Rare in Hawai'i. Found worldwide along continental shelves, bays, estuaries and lagoons of tropical, subtropical, and temperate waters.	43 FR 32800, 76 FR 58868, Dodd 1990, Balazs 1979
Olive Ridley Sea Turtle	<i>Lepidochelys olivacea</i>	Threatened (Entire species, except for the breeding population on the Pacific coast of Mexico, which is listed as endangered)	N/A	Rare in Hawai'i. Occurs worldwide in tropical and warm temperate ocean waters.	43 FR 32800, Pitman 1990, Balacz 1982
Marine mammals					
Blainville's Beaked Whale	<i>Mesoplodon densirostris</i>	Not Listed	Non-strategic	Found worldwide in tropical and temperate waters	Mead 1989
Blue Whale	<i>Balaenoptera musculus</i>	Endangered	Strategic	Acoustically recorded off of Oahu and Midway Atoll, small number of sightings around Hawai'i. Considered extremely rare, generally occur in winter and summer.	35 FR 18319, Bradford et al. 2013, Northrop et al. 1971, Thompson & Friedl 1982, Stafford et al. 2001
Bottlenose Dolphin	<i>Tursiops truncatus</i>	Not Listed	Non-strategic	Distributed worldwide in tropical and warm-temperate waters. Pelagic stock distinct from island-associated stocks.	Perrin et al. 2009, Martien et al. 2012
Bryde's Whale	<i>Balaenoptera edeni</i>	Not Listed	Unknown	Distributed widely across tropical and warm-temperate Pacific Ocean.	Leatherwood et al. 1982
Common Dolphin	<i>Delphinus delphis</i>	Not Listed	N/A	Found worldwide in temperate and subtropical seas.	Perrin et al. 2009
Cuvier's Beaked Whale	<i>Ziphius cavirostris</i>	Not Listed	Non-strategic	Occur year round in Hawaiian waters.	McSweeney et al. 2007
Dall's Porpoise	<i>Phocoenoides dalli</i>	Not Listed	Non-strategic	Range across the entire north Pacific Ocean.	Hall 1979
Dwarf Sperm Whale	<i>Kogia sima</i>	Not Listed	Non-strategic	Most common in waters between 500 m and 1,000 m in depth. Found worldwide in tropical and warm-temperate waters.	Nagorsen 1985, Baird et al. 2013

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
False Killer Whale	<i>Pseudorca crassidens</i>	Not Listed	Non-strategic	Found worldwide in tropical and warm-temperate waters. Pelagic stock tracked to within 11 km of Hawaiian islands.	Stacey et al. 1994, Baird et al. 2012, Bradford et al. 2015
Fin Whale	<i>Balaenoptera physalus</i>	Endangered	Strategic	Infrequent sightings in Hawai'i waters. Considered rare in Hawai'i, though may migrate into Hawaiian waters during fall/winter based on acoustic recordings.	35 FR 18319, Hamilton et al. 2009, Thompson & Friedl 1982
Fraser's Dolphin	<i>Lagenodelphis hosei</i>	Not Listed	Non-strategic	Found worldwide in tropical waters.	Perrin et al. 2009
Guadalupe Fur Seal	<i>Arctocephalus townsendi</i>	Threatened	Strategic	Rare sightings. Little known about their pelagic distribution. Breed mainly on Isla Guadalupe, Mexico.	50 FR 51252, Gallo-Reynoso et al. 2008, Fleischer 1987
Hawaiian Monk Seal	<i>Neomonachus schauinslandi</i>	Endangered ^a	Strategic	Endemic tropical seal. Occurs throughout the archipelago. MHI population spends some time foraging in federal waters during the day.	41 FR 51611, Baker et al. 2011
Humpback Whale	<i>Megaptera novaeangliae</i>	Delisted Due to Recovery (Hawai'i DPS)	Strategic	Migrate through the archipelago and breed during the winter. Common during winter months, when they are generally found within the 100 m isobath.	35 FR 18319, 81 FR 62259, Childerhouse et al. 2008, Wolman & Jurasz 1976, Herman & Antinaja 1977, Rice & Wolman 1978
Killer Whale	<i>Orcinus orca</i>	Not Listed	Non-strategic	Rare in Hawai'i. Prefer colder waters within 800 km of continents.	Mitchell 1975, Baird et al. 2006
Longman's Beaked Whale	<i>Indopacetus pacificus</i>	Not Listed	Non-strategic	Found in tropical waters from the eastern Pacific westward through the Indian Ocean to the eastern coast of Africa. Rare in Hawai'i.	Dalebout 2003, Baird et al. 2013
Melon-Headed Whale	<i>Peponocephala electra</i>	Not Listed	Non-strategic	Found in tropical and warm-temperate waters worldwide, found primarily in equatorial waters. Uncommon in Hawai'i.	Perryman et al. 1994, Barlow 2006, Bradford et al. 2013
Minke Whale	<i>Balaenoptera</i>	Not Listed	Non-strategic	Occur seasonally around	Barlow 2003,

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
	<i>acutorostrata</i>			Hawai'i	Rankin & Barlow 2005
North Pacific Right Whale	<i>Eubalaena japonica</i>	Endangered ^a	Strategic	Extremely rare in Hawai'i waters	35 FR 18319, 73 FR 12024, Rowntree et al. 1980, Herman et al. 1980
Northern Elephant Seal	<i>Mirounga angustirostris</i>	Not Listed	Non-strategic	Females migrate to central North Pacific to feed on pelagic prey	Le Beouf et al. 2000
Northern Fur Seal	<i>Callorhinus ursinus</i>	Not Listed	Non-strategic	Range across the north Pacific Ocean.	Gelatt et al. 2015
Pacific White-Sided Dolphin	<i>Lagenorhynchus obliquidens</i>	Not Listed	Non-strategic	Endemic to temperate waters of North Pacific Ocean. Occur both on the high seas and along continental margins.	Brownell et al. 1999
Pantropical Spotted Dolphin	<i>Stenella attenuata attenuata</i>	Not Listed	Non-strategic	Common and abundant throughout the Hawaiian archipelago. Pelagic stock occurs outside of insular stock areas (20 km for Oahu and 4-island stocks, 65 km for Hawai'i Island stock)	Baird et al. 2013, Oleson et al. 2013
Pygmy Killer Whale	<i>Feresa attenuata</i>	Not Listed	Non-strategic	Small resident population in Hawaiian waters. Found worldwide in tropical and subtropical waters.	McSweeney et al. 2009, Ross & Leatherwood 1994
Pygmy Sperm Whale	<i>Kogia breviceps</i>	Not Listed	Non-strategic	Found worldwide in tropical and warm-temperate waters.	Caldwell & Caldwell 1989
Risso's Dolphin	<i>Grampus griseus</i>	Not Listed	Non-strategic	Found in tropical to warm-temperate waters worldwide.	Perrin et al. 2009
Rough-Toothed Dolphin	<i>Steno bredanensis</i>	Not Listed	Non-strategic	Found in tropical to warm-temperate waters worldwide. Occasionally found offshore of Hawai'i.	Perrin et al. 2009, Bradford et al. 2013, Barlow 2006, Baird et al. 2013
Sei Whale	<i>Balaenoptera borealis</i>	Endangered	Strategic	Rare in Hawai'i. Generally found in offshore temperate waters.	35 FR 18319, Barlow 2003, Bradford et al. 2013
Short-Finned Pilot Whale	<i>Globicephala macrorhynchus</i>	Not Listed	Non-strategic	Found in tropical to warm-temperate waters worldwide. Commonly observed around MHI and present around NWHI.	Shallenberger 1981, Baird et al. 2013, Bradford et al. 2013

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
Sperm Whale	<i>Physeter macrocephalus</i>	Endangered	Strategic	Found in tropical to polar waters worldwide, most abundant cetaceans in the region. Sighted off the NWHI and the MHI.	35 FR 18319, Rice 1960, Lee 1993, Barlow 2006, Mobley et al. 2000, Shallenberger 1981
Spinner Dolphin	<i>Stenella longirostris</i>	Not Listed	Non-strategic	Found worldwide in tropical and warm-temperate waters. Pelagic stock found outside of island-associated boundaries (10 nm)	Perrin et al. 2009
Striped Dolphin	<i>Stenella coeruleoalba</i>	Not Listed	Non-strategic	Found in tropical to warm-temperate waters throughout the world	Perrin et al. 2009
Elasmobranchs					
Giant manta ray	<i>Manta birostris</i>	Threatened	N/A	Found worldwide in tropical, subtropical, and temperate waters. Commonly found in upwelling zones, oceanic island groups, offshore pinnacles and seamounts, and on shallow reefs.	Dewar et al. 2008, Marshall et al. 2009, Marshall et al. 2011.
Oceanic whitetip shark	<i>Carcharhinus longimanus</i>	Threatened	N/A	Found worldwide in open ocean waters from the surface to 152 m depth. It is most commonly found in waters > 20°C	Bonfil et al. 2008, Backus et al, 1956, Strasburg 1958, Compagno 1984
Scalloped hammerhead shark	<i>Sphyrna lewini</i>	Endangered (Eastern Pacific DPS)	N/A	Found in coastal areas from southern California to Peru.	Compagno 1984, Baum et al. 2007, Bester 2011
Scalloped hammerhead shark	<i>Sphyrna lewini</i>	Threatened (Indo-West Pacific DPS)	N/A	Occur over continental and insular shelves, and adjacent deep waters, but rarely found in waters < 22°C. Range from the intertidal and surface to depths up to 450–512 m.	Compagno 1984, Schulze-Haugen & Kohler 2003, Sanches 1991, Klimley 1993
Corals					
N/A	<i>Acropora globiceps</i>	Threatened	N/A	Occur on upper reef slopes, reef flats, and adjacent habitats in depths ranging from 0 to 8 m.	Veron 2014

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
N/A	<i>Acropora jacquelineae</i>	Threatened	N/A	Found in numerous subtidal reef slope and back-reef habitats, including but not limited to, lower reef slopes, walls and ledges, mid-slopes, and upper reef slopes protected from wave action, and depth range is 10 to 35 m.	Veron 2014
N/A	<i>Acropora retusa</i>	Threatened	N/A	Occur in shallow reef slope and back-reef areas, such as upper reef slopes, reef flats, and shallow lagoons, and depth range is 1 to 5 m.	Veron 2014
N/A	<i>Acropora speciosa</i>	Threatened	N/A	Found in protected environments with clear water and high diversity of <i>Acropora</i> and steep slopes or deep, shaded waters. Depth range is 12 to 40 meters, and it has been found in mesophotic habitat (40-150 m).	Veron 2014
N/A	<i>Euphyllia paradivisa</i>	Threatened	N/A	Found in environments protected from wave action on at least upper reef slopes, mid-slope terraces, and lagoons in depths ranging from 2 to 25 m depth.	Veron 2014
N/A	<i>Isopora crateriformis</i>	Threatened	N/A	Found in shallow, high-wave energy environments, from low tide to at least 12 m deep, and have been reported from mesophotic depths (less than 50 m depth).	Veron 2014
N/A	<i>Seriatopora aculeata</i>	Threatened	N/A	Found in broad range of habitats including, but not limited to, upper reef slopes, mid-slope terraces, lower reef slopes, reef flats, and lagoons, and depth ranges from 3 to 40 m.	Veron 2014
Invertebrates					

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
Chambered nautilus	<i>Nautilus pompilius</i>	Threatened	N/A	Found in small, isolated populations throughout the Indo-Pacific on steep-sloped forereefs with sandy, silty, or muddy bottom substrates from depths of 100 m to 500 m.	83 FR 48948, CITES 2016

^a These species have critical habitat designated under the ESA. See Table B-4 .

Table B-3. Protected species found or reasonably believed to be found near or in American Samoa longline waters.

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
Seabirds					
Audubon's Shearwater	<i>Puffinus lherminieri</i>	Not Listed	N/A	Resident	Craig 2005
Black Noddy	<i>Anous minutus</i>	Not Listed	N/A	Resident	Craig 2005
Black-Naped Tern	<i>Sterna sumatrana</i>	Not Listed	N/A	Visitor	Craig 2005
Blue-Gray Noddy	<i>Procelsterna cerulea</i>	Not Listed	N/A	Resident	Craig 2005
Bridled Tern	<i>Onychoprion anaethetus</i>	Not Listed	N/A	Visitor	Craig 2005
Brown Booby	<i>Sula leucogaster</i>	Not Listed	N/A	Resident	Craig 2005
Brown Noddy	<i>Anous stolidus</i>	Not Listed	N/A	Resident	Craig 2005
Christmas Shearwater	<i>Puffinus nativitatis</i>	Not Listed	N/A	Resident?	Craig 2005
Collared Petrel	<i>Pterodroma brevipes</i>	Not Listed	N/A	Resident?	Craig 2005
White Tern	<i>Gygis alba</i>	Not Listed	N/A	Resident	Craig 2005
Greater Crested Tern	<i>Thalasseus bergii</i>	Not Listed	N/A	Visitor	Craig 2005
Gray-Backed Tern	<i>Onychoprion lunatus</i>	Not Listed	N/A	Resident	Craig 2005
Great Frigatebird	<i>Fregata minor</i>	Not Listed	N/A	Resident	Craig 2005
Herald Petrel	<i>Pterodroma heraldica</i>	Not Listed	N/A	Resident	Craig 2005
Laughing Gull	<i>Leucophaeus atricilla</i>	Not Listed	N/A	Visitor	Craig 2005
Lesser Frigatebird	<i>Fregata ariel</i>	Not Listed	N/A	Resident	Craig 2005
Masked Booby	<i>Sula dactylatra</i>	Not Listed	N/A	Resident	Craig 2005
Newell's Shearwater	<i>Puffinus auricularis newelli</i>	Threatened	N/A	Visitor	40 FR 44149, Craig 2005
Red-Footed Booby	<i>Sula sula</i>	Not Listed	N/A	Resident	Craig 2005

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
Red-Tailed Tropicbird	<i>Phaethon rubricauda</i>	Not Listed	N/A	Resident	Craig 2005
Short-Tailed Shearwater	<i>Ardenna tenuirostris</i>	Not Listed	N/A	Visitor	Craig 2005
Sooty Shearwater	<i>Ardenna grisea</i>	Not Listed	N/A	Visitor	Craig 2005
Sooty Tern	<i>Sterna fuscata</i>	Not Listed	N/A	Resident	Craig 2005
Tahiti Petrel	<i>Pterodroma rostrata</i>	Not Listed	N/A	Resident	Craig 2005
Wedge-Tailed Shearwater	<i>Ardenna pacifica</i>	Not Listed	N/A	Resident?	Craig 2005
White-Necked Petrel	<i>Pterodroma cervicalis</i>	Not Listed	N/A	Visitor	Craig 2005
White-Faced Storm-Petrel	<i>Pelagodroma marina</i>	Not Listed	N/A	Visitor	Craig 2005
White-Tailed Tropicbird	<i>Phaethon lepturus</i>	Not Listed	N/A	Resident	Craig 2005
White-Throated Storm-Petrel	<i>Nesofregatta fuliginosa</i>	Not Listed	N/A	Resident?	Craig 2005
Laysan Albatross	<i>Phoebastria immutabilis</i>	Not Listed	N/A	Breed mainly in Hawai'i, and range across the North Pacific Ocean.	Causey 2008
Hawaiian Petrel	<i>Pterodroma sandwichensis</i> (<i>Pterodroma phaeopygia sandwichensis</i>)	Endangered	N/A	Breed in MHI, and range across the central Pacific Ocean.	32 FR 4001, Simons & Hodges 1998
Laysan Albatross	<i>Phoebastria immutabilis</i>	Not Listed	N/A	Breed mainly in Hawai'i, and range across the North Pacific Ocean.	Causey 2009
Northern Fulmar	<i>Fulmarus glacialis</i>	Not Listed	N/A	Breed and range across North Pacific Ocean.	Hatch & Nettleship 2012
Short-Tailed Albatross	<i>Phoebastria albatrus</i>	Endangered	N/A	Breed in Japan and NWHI, and range across the North Pacific Ocean.	35 FR 8495, 65 FR 46643, BirdLife International 2017
Sea turtles					
Green Sea Turtle	<i>Chelonia mydas</i>	Endangered (Central South Pacific DPS)	N/A	Frequently seen. Nest at Rose Atoll in small numbers.	43 FR 32800, 81 FR 20057, Balacz 1994
Hawksbill Sea Turtle	<i>Eretmochelys imbricata</i>	Endangered ^a	N/A	Frequently seen. Nest at Rose Atoll, Swain's Island, and Tutuila.	35 FR 8491, NMFS & USFWS 2013, Tuato'o-Bartley et al. 1993
Leatherback Sea Turtle	<i>Dermochelys coriacea</i>	Endangered ^a	N/A	Very rare. One juvenile recovered dead in experimental longline fishing.	35 FR 8491, Grant 1994

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
Loggerhead Sea Turtle	<i>Caretta caretta</i>	Endangered (South Pacific DPS)	N/A	No known sightings. Found worldwide along continental shelves, bays, estuaries and lagoons of tropical, subtropical, and temperate waters.	43 FR 32800, 76 FR 58868, Utzurrum 2002, Dodd 1990
Olive Ridley Sea Turtle	<i>Lepidochelys olivacea</i>	Threatened (Entire species, except for the endangered breeding population on the Pacific coast of Mexico)	N/A	Rare. Three known sightings.	43 FR 32800, Utzurrum 2002
Marine mammals					
Blainville's Beaked Whale	<i>Mesoplodon densirostris</i>	Not Listed	Non-strategic	Found worldwide in tropical and temperate waters	Mead 1989
Blue Whale	<i>Balaenoptera musculus</i>	Endangered	Strategic	No known sightings. Occur worldwide, and are known to be found in the western South Pacific.	35 FR 18319, Olson et al. 2015
Bottlenose Dolphin	<i>Tursiops truncatus</i>	Not Listed	Non-strategic	Distributed worldwide in tropical and warm-temperate waters. Pelagic stock distinct from island-associated stocks.	Perrin et al. 2009, Martien et al. 2012
Bryde's Whale	<i>Balaenoptera edeni</i>	Not Listed	Unknown	Distributed widely across tropical and warm-temperate Pacific Ocean.	Leatherwood et al. 1982
Common Dolphin	<i>Delphinus delphis</i>	Not Listed	N/A	Found worldwide in temperate and subtropical seas.	Perrin et al. 2009
Cuvier's Beaked Whale	<i>Ziphius cavirostris</i>	Not Listed	Non-strategic	Occur worldwide.	Heyning 1989
Dwarf Sperm Whale	<i>Kogia sima</i>	Not Listed	Non-strategic	Found worldwide in tropical and warm-temperate waters.	Nagorsen 1985
False Killer Whale	<i>Pseudorca crassidens</i>	Not Listed	Unknown	Found in waters within the U.S. EEZ of A. Samoa	Bradford et al. 2015
Fin Whale	<i>Balaenoptera physalus</i>	Endangered	Strategic	No known sightings but reasonably expected to occur in A. Samoa. Found worldwide.	35 FR 18319, Hamilton et al. 2009
Fraser's Dolphin	<i>Lagenodelphis hosei</i>	Not Listed	Non-strategic	Found worldwide in tropical waters.	Perrin et al. 2009

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
Guadalupe Fur Seal	<i>Arctocephalus townsendi</i>	Threatened	Strategic	No known sightings. Little known about their pelagic distribution. Breed mainly on Isla Guadalupe, Mexico.	50 FR 51252, Gallo-Reynoso et al. 2008, Fleischer 1987
Humpback Whale	<i>Megaptera novaeangliae</i>	Delisted Due to Recovery (Oceania DPS)	Strategic	Migrate through the archipelago and breed during the winter in American Samoan waters.	35 FR 18319, 81 FR 62259,, Guarrige et al. 2007, SPWRC 2008
Killer Whale	<i>Orcinus orca</i>	Not Listed	Non-strategic	Found worldwide. Prefer colder waters within 800 km of continents.	Leatherwood & Dalheim 1978, Mitchell 1975, Baird et al. 2006
Longman's Beaked Whale	<i>Indopacetus pacificus</i>	Not Listed	Non-strategic	Found in tropical waters from the eastern Pacific westward through the Indian Ocean to the eastern coast of Africa.	Dalebout 2003
Melon-Headed Whale	<i>Peponocephala electra</i>	Not Listed	Non-strategic	Found in tropical and warm-temperate waters worldwide, primarily found in equatorial waters.	Perryman et al. 1994
Minke Whale	<i>Balaenoptera acutorostrata</i>	Not Listed	Non-strategic	Uncommon in this region, usually seen over continental shelves in the Pacific Ocean.	Brueggeman et al. 1990
North Pacific Right Whale	<i>Eubalaena japonica</i>	Endangered ^a	Strategic	Extremely rare.	35 FR 18319, 73 FR 12024, Childerhouse et al. 2008, Wolman & Jurasz 1976, Herman & Antinaja 1977, Rice & Wolman 1978
Northern Elephant Seal	<i>Mirounga angustirostris</i>	Not Listed	Non-strategic	Females migrate to central North Pacific to feed on pelagic prey	Le Beouf et al. 2000
Pantropical Spotted Dolphin	<i>Stenella attenuata attenuata</i>	Not Listed	Non-strategic	Found in tropical and subtropical waters worldwide.	Perrin et al. 2009
Pygmy Killer Whale	<i>Feresa attenuata</i>	Not Listed	Non-strategic	Found in tropical and subtropical waters worldwide.	Ross & Leatherwood 1994
Pygmy Sperm Whale	<i>Kogia breviceps</i>	Not Listed	Non-strategic	Found worldwide in tropical and warm-temperate waters.	Caldwell & Caldwell 1989

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
Risso's Dolphin	<i>Grampus griseus</i>	Not Listed	Non-strategic	Found in tropical to warm-temperate waters worldwide.	Perrin et al. 2009
Rough-Toothed Dolphin	<i>Steno bredanensis</i>	Not Listed	Unknown	Found in tropical to warm-temperate waters worldwide. Common in A. Samoa waters.	Perrin et al. 2009, Craig 2005
Sei Whale	<i>Balaenoptera borealis</i>	Endangered	Strategic	Generally found in offshore temperate waters.	35 FR 18319, Barlow 2003, Bradford et al. 2013
Short-Finned Pilot Whale	<i>Globicephala macrorhynchus</i>	Not Listed	Non-strategic	Found in tropical to warm-temperate waters worldwide	Shallenberger 1981, Baird et al. 2013, Bradford et al. 2013
Sperm Whale	<i>Physeter macrocephalus</i>	Endangered	Strategic	Found in tropical to polar waters worldwide, most abundant cetaceans in the region.	35 FR 18319, Rice 1960, Barlow 2006, Lee 1993, Mobley et al. 2000, Shallenberger 1981
Spinner Dolphin	<i>Stenella longirostris</i>	Not Listed	Unknown	Common in American Samoa, found in waters with mean depth of 44 m.	Reeves et al. 1999, Johnston et al. 2008
Striped Dolphin	<i>Stenella coeruleoalba</i>	Not Listed	Non-strategic	Found in tropical to warm-temperate waters throughout the world	Perrin et al. 2009
Elasmobranchs					
Giant manta ray	<i>Manta birostris</i>	Threatened	N/A	Found worldwide in tropical, subtropical, and temperate waters. Commonly found in upwelling zones, oceanic island groups, offshore pinnacles and seamounts, and on shallow reefs.	Dewar et al. 2008, Marshall et al. 2009, Marshall et al. 2011.
Oceanic whitetip shark	<i>Carcharhinus longimanus</i>	Threatened	N/A	Found worldwide in open ocean waters from the surface to 152 m depth. It is most commonly found in waters > 20°C.	Bonfil et al. 2008, Backus et al, 1956, Strasburg 1958, Compagno 1984
Scalloped hammerhead shark	<i>Sphyrna lewini</i>	Threatened (Indo-West Pacific DPS)	N/A	Occur over continental and insular shelves, and adjacent deep waters, but rarely found in waters < 22°C. Range from the intertidal and surface to depths up to 450–512 m.	Compagno 1984, Schulze-Haugen & Kohler 2003, Sanches 1991, Klimley 1993

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
Corals					
N/A	<i>Acropora globiceps</i>	Threatened	N/A	Occur on upper reef slopes, reef flats, and adjacent habitats in depths from 0 to 8 m	Veron 2014
N/A	<i>Acropora jacquelineae</i>	Threatened	N/A	Found in numerous subtidal reef slope and back-reef habitats, including but not limited to, lower reef slopes, walls and ledges, mid-slopes, and upper reef slopes protected from wave action, and its depth range is 10 to 35 m.	Veron 2014
N/A	<i>Acropora retusa</i>	Threatened	N/A	Occur in shallow reef slope and back-reef areas, such as upper reef slopes, reef flats, and shallow lagoons. Depth range is 1 to 5 m.	Veron 2014
N/A	<i>Acropora speciosa</i>	Threatened	N/A	Found in protected environments with clear water and high diversity of Acropora and steep slopes or deep, shaded waters. Depth range is 12 to 40 meters, and have been found in mesophotic habitat (40-150 m).	Veron 2014
N/A	<i>Euphyllia paradivisa</i>	Threatened	N/A	Found in environments protected from wave action on at least upper reef slopes, mid-slope terraces, and lagoons in depths ranging from 2 to 25 m depth.	Veron 2014
N/A	<i>Isopora crateriformis</i>	Threatened	N/A	Found in shallow, high-wave energy environments, from low tide to at least 12 meters deep, and have been reported from mesophotic depths (less than 50 m depth).	Veron 2014
Invertebrates					

Common name	Scientific name	ESA listing status	MMPA status	Occurrence	References
Chambered nautilus	<i>Nautilus pompilius</i>	Threatened	N/A	Found in small, isolated populations throughout the Indo-Pacific on steep-sloped forereefs with sandy, silty, or muddy bottom substrates from depths of 100 m to 500 m.	83 FR 48948, CITES 2016

^a These species have critical habitat designated under the ESA. See Table B-4.

Table B-4. ESA-listed species' critical habitat in the Pacific Ocean^a.

Common Name	Scientific Name	ESA Listing Status	Critical Habitat	References
Hawksbill Sea Turtle	<i>Eretmochelys imbricata</i>	Endangered	None in the Pacific Ocean.	63 FR 46693
Leatherback Sea Turtle	<i>Dermochelys coriacea</i>	Endangered	Approximately 16,910 square miles (43,798 square km) stretching along the California coast from Point Arena to Point Arguello east of the 3,000 meter depth contour; and 25,004 square miles (64,760 square km) stretching from Cape Flattery, Washington to Cape Blanco, Oregon east of the 2,000 meter depth contour.	77 FR 4170
Hawaiian Monk Seal	<i>Neomonachus schauinslandi</i>	Endangered	Ten areas in the Northwestern Hawaiian Islands (NWHI) and six in the main Hawaiian Islands (MHI). These areas contain one or a combination of habitat types: Preferred pupping and nursing areas, significant haul-out areas, and/or marine foraging areas, that will support conservation for the species.	53 FR 18988, 51 FR 16047, 80 FR 50925
North Pacific Right Whale	<i>Eubalaena japonica</i>	Endangered	Two specific areas are designated, one in the Gulf of Alaska and another in the Bering Sea, comprising a total of approximately 95,200 square kilometers (36,750 square miles) of marine habitat.	73 FR 19000, 71 FR 38277

^a For maps of critical habitat, see <https://www.fisheries.noaa.gov/national/endangered-species-conservation/critical-habitat>.

REFERENCES

Andrews, K.R., Karczmarski, L., Au, W.W.L., Rickards, S.H., Vanderlip, C.A., Bowen, B.W., Grau, E.G., and Toonen, R.J., 2010. Rolling stones and stable homes: social structure, habitat diversity and population genetics of the Hawaiian spinner dolphin (*Stenella longirostris*). *Molecular Ecology*, 19, pp. 732-748.

- Awkerman, J.A., Anderson, D.J., and Whittow, G.C., 2009. Laysan Albatross (*Phoebastria immutabilis*). In: Rodewald, P.G. [ed.]. The Birds of North America. Ithaca: Cornell Lab of Ornithology. Retrieved from the Birds of North America at <https://birdsna.org/Species-Account/bna/species/layalb>.
- Backus, R.H., Springer, S., and Arnold, E.L., 1956. A contribution to the natural history of the white tip shark, *Pseudocarcharias kamoharui* (Poey). *Deep-Sea Research*, 3, pp. 178-188.
- Baird, R.W., McSweeney, D.J., Bane, C., Barlow, J., Salden, D.R., Antoine, L.R.K., LeDuc, R.G. and Webster, D.L., 2006. Killer Whales in Hawaiian Waters: Information on Population Identity and Feeding Habits. *Pacific Science*, 60(4), pp. 523-530.
- Baird, R.W., Gorgone, A.M., McSweeney, D.J., Ligon, A.D., Deakos, M.H., Webster, D.L., Schorr, G.S., Martien, K.K., Salden, D.R., and Mahaffy, S.D., 2009. Population structure of island-associated dolphins: Evidence from photo-identification of common bottlenose dolphins (*Tursiops truncatus*) in the main Hawaiian Islands. *Marine Mammal Science*, 25, pp. 251-274.
- Baird, R.W., Webster, D.L., Aschettino, J.M., Schorr, G.S. and McSweeney, D.J., 2013. Odontocete cetaceans around the main Hawaiian Islands: Habitat use and relative abundance from small-boat sighting surveys. *Aquatic Mammals*, 39(3), 253 pp.
- Baker J.D., Harting, A.L., Wurth, T.A., and Johanos, T.C., 2011. Dramatic shifts in Hawaiian monk seal distribution predicted from divergent regional trends. *Marine Mammal Science*, 27(1), pp. 78-93.
- Balazs, G.H. 1979. Loggerhead turtle recovered from a tiger shark at Kure Atoll. *'Elepaio*, 39(12), pp. 45-47.
- Balazs, G.H. 1982. Status of sea turtles in the central Pacific Ocean. In: Bjorndal, K.A. [ed.]. Biology and Conservation of Sea Turtles. *Smithsonian Institution Scholarly Press*, Washington, D.C., 583 pp.
- Balazs, G.H., Craig, P., Winton, B.R. and Miya, R.K., 1994. Satellite telemetry of green turtles nesting at French Frigate Shoals, Hawaii, and Rose Atoll, American Samoa. In: Bjorndal, K.A., Bolten, A.B., Johnson, D.A., and Eliazar, P.J. [eds.]. Proceedings of the fourteenth annual symposium on sea turtle biology and conservation.
- Balazs, G.H., Hirth, H., Kawamoto, P., Nitta, E., Ogren, L., Wass, R., and Wetherall, J., 1992. Interim Recovery Plan for Hawaiian Sea Turtles. Honolulu Lab, Southwest Fisheries Science Center, National Marine Fisheries Service, NOAA, Honolulu, HI 96822-2396. Southwest Fisheries Science Center Administrative Report H-92-01. 76 pp.
- Barlow, J., 2003. Preliminary Estimates of the Abundance of Cetaceans along the U.S. West Coast, 1991-2001. U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Southwest Fisheries Science Center.

- Barlow, J., 2006. Cetacean abundance in Hawaiian waters estimated from a summer/fall survey in 2002. *Marine Mammal Science*, 22(2), pp. 446-464.
- Baum, J., Clarke, S., Domingo, A., Ducrocq, M., Lamónaca, A.F., Gaibor, N., Graham, R., Jorgensen, S., Kotas, J.E., Medina, E., Martinez-Ortiz, J., Monzini Taccone di Sitizano, J., Morales, M.R., Navarro, S.S., Pérez-Jiménez, J.C., Ruiz, C., Smith, W., Valenti, S.V. and Vooren, C.M., 2007. *Sphyrna lewini*. The IUCN Red List of Threatened Species 2007: e.T39385A10190088. Downloaded on 21 Feb 2017.
- Baum, J.K., Medina, E., Musick, J.A., and Smale, M., 2006. *Carcharhinus longimanus*. In: IUCN 2007. 2007 IUCN Red List of Threatened Species.
- Bester, C., 2011. Species Profile: Scalloped Hammerhead. Florida Museum of Natural History. <http://www.flmnh.ufl.edu/fish/Gallery/Descript/Schammer/ScallopedHammerhead.html>.
- BirdLife International, 2017. Species factsheet: *Phoebastria albatrus*. Downloaded from <http://www.birdlife.org> on 02/04/2017.
- Bonfil, R., Clarke, S., Nakano, H., Camhi, M.D., Pikitch, E.K. and Babcock, E.A., 2008. The biology and ecology of the oceanic whitetip shark, *Carcharhinus longimanus*. In: Camhi, M.D., Pikitch, E.K., and Babcock, E.A. [eds.]. *Sharks of the Open Ocean: Biology, Fisheries, and Conservation*, pp.128-139.
- Bradford, A.L., Oleson, E.M., Baird, R.W., Boggs, C.H., Forney, K.A., and Young, N.C., 2015. Revised stock boundaries for false killer whales (*Pseudorca crassidens*) in Hawaiian waters. NOAA Tech. Memo. NMFS-PIFSC-47.
- Bradford, A.L., K.A. Forney, E.M. Oleson, and J. Barlow. 2013. Line-transect abundance estimates of cetaceans in the Hawaiian EEZ. PIFSC Working Paper WP-13-004.
- Brownell, J. R. L., W. A. Walker, and K. A. Forney. 1999. Pacific white-sided dolphin, *Lagenorhynchus obliquidens* Gill, 1865. In: Ridgeway, S.H., and Harrison, R. [eds.]. *Handbook of marine mammals: the second book of dolphins and the porpoises*. Academic Press, San Diego, CA.
- Brueggeman, J.J., Green, G.A., Balcomb, K.C., Bowlby, C.E., Grotiefendt, R.A., Briggs, K.T., Bonnell, M.L., Ford, R.G., Varoujean, D.H., Heinemann, D. and Chapman D.G., 1990. Oregon-Washington Marine Mammal and Seabird Survey: Information synthesis and hypothesis formulation. U.S. Department of the Interior, OCS Study MMS 89-0030.
- Caldwell, D.K. and Caldwell, M.C., 1989. Pygmy sperm whale *Kogia breviceps* (de Blainville, 1838)/dwarf sperm whale *Kogia simus* (Owen, 1866). *Handbook of marine mammals*, 4, pp. 235-260.
- Childerhouse, S., J. Jackson, C. S. Baker, N. Gales, P. J. Clapham, and R. L. Brownell, Jr., 2008. *Megaptera novaeangliae*, Oceania subpopulation. IUCN Red List of Threatened Species <http://www.iucnredlist.org/details/132832>.

- CITES (2016) Consideration of proposals for amendment of Appendices I and II: Nautilidae Convention on International Trade in Endangered Species of Wild Fauna and Flora; Seventeenth meeting of the Conference of the Parties; 24 September – 5 October 2016, Johannesburg, South Africa
- Cliffton, K., Cornejo, D.O., and Felger, R.S., 1982. Sea turtles of the Pacific coast of Mexico. *In*: Bjorndal, K.A. [ed.]. *Biology and Conservation of Sea Turtles*. Smithsonian Inst. Press, Washington, D.C., 583 pp.
- Compagno, L.J.V., 1984. *FAO Species Catalogue*. Vol. 4. *Sharks of the World. An Annotated and Illustrated Catalogue of Shark Species Known to Date. Carcharhiniformes*. FAO Fish Synop 124, Vol. 4, Part 2.
- Craig, P., 2005. *Natural history guide to American Samoa*. National Park of American Samoa, Department of Marine and Wildlife Resources, American Samoa Community College.
- Dalebout, M.L., Baker, C.S., Anderson, R.C., Best, P.B., Cockcroft, V.G., Hinsz, H.L., Peddemors, V. and Pitman, R.L., 2003. Appearance, distribution, and genetic distinctiveness of Longman's beaked whale, *Indopacetus pacificus*. *Marine Mammal Science*, 19(3), pp. 421-461.
- Dewar, H., Mous, P., Domeier, M., Muljadi, A., Pet, J., and Whitty, J., 2008. Movements and site fidelity of the giant manta ray, *Manta birostris*, in the Komodo Marine Park, Indonesia. *Marine Biology*, 155, pp. 121-133.
- Dodd, C.K., 1990. *Caretta caretta* (Linnaeus) Loggerhead Sea Turtle. *Catalogue of American Amphibians and Reptiles*, pp. 483.1-483.7.
- Eckert, K.L., B.P. Wallace, J.G. Frazier, S.A. Eckert, and P.C.H. Pritchard. 2012. Synopsis of the biological data on the leatherback sea turtle (*Dermochelys coriacea*). U.S. Department of Interior, Fish and Wildlife Service, Biological Technical Publication BTP-R4015-2012, Washington, D.C.
- Fleischer, L.A., 1987. Guadalupe fur seal, *Arctocephalus townsendi*. *In*: Croxall, J.P. and Gentry R.L. [eds.]. *Status, biology, and ecology of fur seals*. National Oceanic and Atmospheric Association, Technical Report, National Marine Fisheries Service 51, pp. 1-212.
- Gallo-Reynoso, J.P., Figueroa-Carranza, A.L. and Le Boeuf, B.J., 2008. Foraging behavior of lactating Guadalupe fur seal females. *In*: Lorenzo, C., Espinoza, E., and Ortega, J. [eds.]. *Avances en el Estudio de los Mamíferos de México. Publicaciones Especiales*, 2, pp. 595-614.
- Garrigue, C., Franklin, T., Russell, K., Burns, D., Poole, M., Paton, D., Hauser, N., Oremus, M., Constantine, R., Childerhouse, S., Mattila, D., Gibbs, N., Franklin, W., Robbins, J., Clapham, P., and Baker, C.S., 2007. First assessment of interchange of humpback whales between Oceania and the east coast of Australia. International Whaling Commission, Anchorage, Alaska. SC/59/SH15.

- Gelatt, T., Ream, R., and Johnson, D., 2015. *Callorhinus ursinus*. The IUCN Red List of Threatened Species 2015:e.T3590A45224953. Available at <http://www.iucnredlist.org/details/3590/0>.
- Grant, G.S. 1994. Juvenile leatherback turtle caught by longline fishing in American Samoa. *Marine Turtle Newsletter*, 66, pp. 3-5.
- Hall, J., 1979. A survey of cetaceans of Prince William Sound and adjacent waters - their numbers and seasonal movements. Unpubl. rep. to Alaska Outer Continental Shelf Environmental Assessment Programs. NOAA OCSEAP Juneau Project Office, Juneau, AK. 37 pp.
- Hamilton, T.A., Redfern, J.V., Barlow, J., Balance, L.T., Gerrodette, T., Holt, R.S., Forney, K.A., and Taylor, B.L., 2009. Atlas of cetacean sightings for Southwest Fisheries Science Center Cetacean and Ecosystem Surveys: 1986-2005. U.S. Dep. of Commerce, NOAA Technical Memorandum, NOAA-TM-NMFSSWFSC-440. 70 pp.
- Herman, L.M. and Antinaja, R.C., 1977. Humpback whales in the Hawaiian breeding waters: Population and pod characteristics. *Scientific Reports of the Whales Research Institute*, 29, pp. 59-85.
- Herman, L. M., C. S. Baker, P. H. Forestell, and R. C. Antinaja. 1980. Right whale, *Balaena glacialis*, sightings near Hawaii: a clue to the wintering grounds? *Marine Ecology Progress Series*, 2, pp. 271-275.
- Heyning, J.E. 1989. Cuvier's beaked whale, *Ziphius cavirostris* (Cuvier, 1823). pp. 289-308. In: S.H. Ridgway and R. Harrison (eds.) Handbook of Marine Mammals. Vol. 4. River Dolphins and the Larger Toothed Whales. *Academic Press*, London and San Diego. 442 pp.
- Hill, M.C., Oleson, E.M., and Andrews, K.R., 2010. New island-associated stocks for Hawaiian spinner dolphins (*Stenella longirostris longirostris*): Rationale and new stock boundaries. Pacific Islands Fisheries Science Center Admin Report H-10-04, 12 pp.
- Johnston, D.W., Robbins, J., Chapla, M.E., Mattila, D.K. and Andrews, K.R., 2008. Diversity, Johnston, D.W., Robbins, J., Chapla, M.E., Mattila, D.K. and Andrews, K.R., 2008. Diversity, habitat associations and stock structure of odontocete cetaceans in the waters of American Samoa, 2003-2006. *Journal of Cetacean Research and Management*, 10(1), pp. 59-66.
- Karczmarski, L., Würsig, B., Gailey, G., Larson, K.W., and Vanderlip, C., 2005. Spinner dolphins in a remote Hawaiian atoll: social grouping and population structure. *Behavioral Ecology*, 16, pp. 675-685.
- Karl, S.A. and Bowen, B.W., 1999. Evolutionary significant units versus geopolitical taxonomy: molecular systematics of an endangered sea turtle (genus *Chelonia*). *Conservation Biology*, 13, pp. 990-999.

- Katahira, L.K., C.M. Forbes, A.H. Kikuta, G.H. Balazs, and M. Bingham. 1994. Recent findings and management of hawksbill turtle nesting beaches in Hawaii. *In*: Bjorndal, K.A., Bolton, A.B., Johnson, D.A., and Eliazar, P.J. [eds.], Proc. of the Fourteenth Annual Symposium on Sea Turtle Biology and Conservation. NOAA Tech. Memo. NMFS-SEFSC-351, 323 pp.
- Klimley, A.P., 1993. Highly directional swimming by scalloped hammerhead sharks, *Sphyrna lewini*, and subsurface irradiance, temperature, bathymetry, and geomagnetic field. *Marine Biology*, 117(1), pp. 1-22.
- Kolinski, S.P., Parker, D.M., Ilo, L.I., and Ruak, J.K., 2001. An assessment of sea turtles and their marine and terrestrial habitats at Saipan, Commonwealth of the Northern Mariana Islands. *Micronesica*, 34, pp. 55-72.
- Le Boeuf, B.J., Crocker, D.E., Costa, D.P., Blackwell, S.B., Webb, P.M. and Houser, D.S. 2000. Foraging ecology of northern elephant seals. *Ecological Monographs*, 70(3), pp. 353-382.
- Lee, T. 1993. Summary of cetacean survey data collected between the years of 1974 and 1985. NOAA Tech.Mem. NMFS 181, 184 pp.
- Leatherwood, J.S. and Dahlheim, M.E., 1978. Worldwide distribution of pilot whales and killer whales. Naval Ocean System Center Technical Report 443, pp. 1-39.
- Leatherwood, S., Reeves, R.R., Perrin, W.F., and Evans, W.E. 1982. Whales, dolphins, and porpoises of the North Pacific and adjacent arctic waters. NOAA Tech. Rep. NMFS Circ. No. 444.
- Mallory M.L., Hatch, S.A., and Nettleship, D.N. 2012. Northern Fulmar (*Fulmarus glacialis*). *In*: Rodewald, P.G. [ed.]. The Birds of North America. Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America at <https://birdsna.org/Species-Account/bna/species/norful>.
- Marshall, A., L.J.V. Compagno, and M.B. Bennett. 2009. Redescription of the genus *Manta* with resurrection of *Manta alfredi* (Krefft, 1868) (Chondrichthyes; Myliobatoidei; Mobulidae). *Zootaxa*, 2301, pp. 1-28.
- Marshall, A., Bennett, M.B., Kodja, G., Hinojosa-Alvarez, S., Galvan-Magana, F., Harding, M., Stevens, G., and Kashiwagi, T., 2011. *Manta birostris*. The IUCN Red List of Threatened Species 2011: e.T198921A9108067.
- Martien, K.K., Baird, R.W., Hedrick, N.M., Gorgone, A.M., Thieleking, J.L., McSweeney, D.J., Robertson, K.M. and Webster, D.L. 2012. Population structure of island-associated dolphins: Evidence from mitochondrial and microsatellite markers for common bottlenose dolphins (*Tursiops truncatus*) around the main Hawaiian Islands. *Marine Mammal Science*, 25(3), pp. 208-232.

- McSweeney, D.J., Baird, R.W., and Mahaffy, S.D., 2007. Site fidelity, associations and movements of Cuvier's (*Ziphius cavirostris*) and Blainville's (*Mesoplodon densirostris*) beaked whales off the island of Hawai'i. *Marine Mammal Science*, 23, pp. 666-687.
- McSweeney, D.J., Baird, R.W., Mahaffy, S.D., Webster, D.L., and Schorr, G.S. 2009. Site fidelity and association patterns of a rare species: Pygmy killer whales (*Feresa attenuata*) in the main Hawaiian Islands. *Marine Mammal Science*, 25, pp. 557-572.
- Mead, J.G., 1989. Beaked whales of the genus *Mesoplodon*. In: Ridgeway, S.H. & Harrison, R. [eds.]. Handbook of marine mammals. Volume 4. River dolphins and the larger toothed whales. Academic Press Ltd: London, 452 pp.
- Mitchell, E., 1975. Report on the meeting on smaller cetaceans, Montreal, April 1-11, 1974. *Journal of the Fisheries Research Board of Canada*, 32(7), pp. 914-916
- Mobley, J.R. Jr., Spitz, S.S., Forney, K.A., Grotenfendt, R., and Forestell, P.H., 2000. Distribution and abundance of odontocete species in Hawaiian waters: preliminary results of 1993-98 aerial surveys. Southwest Fisheries Science Center Administrative Report LJ-00-14C. La Jolla, CA 92037. 26 pp.
- Nagorsen, D., 1985. *Kogia simus*. *Mammalian Species*, 239, pp. 1-6.
- NMFS and USFWS, 1998. Recovery Plan for U.S. Pacific Populations of the Leatherback Turtle (*Dermochelys coriacea*). National Marine Fisheries Service, Silver Spring, MD. 77 pp.
- NMFS and USFWS, 2007. Hawksbill Sea Turtle (*Eretmochelys imbricata*) 5-Year Review: Summary and Evaluation. National Marine Fisheries Service, Silver Spring, Maryland, and U.S. Fish and Wildlife Service Jacksonville, Florida. 90 pp.
- Norris, K.S. and Dohl, T.P., 1980. Behavior of the Hawaiian spinner dolphin, *Stenella longirostris*. *Fisheries Bulletin*, 77, pp. 821-849.
- Norris, K.S., Würsig, B., Wells, R.S., and Würsig, M., 1994. The Hawaiian Spinner Dolphin. *University of California Press*, 408 pp.
- Northrop, J., Cummings, W.C., and Morrison, M.F. 1971. Underwater 20-Hz signals recorded near Midway Island. *Journal of the Acoustical Society of America*, 49, pp. 1909-1910.
- Oleson, E.M., Boggs, C.H., Forney, K.A., Hanson, M.B., Kobayashi, D.R., Taylor, B.L., Wade, P.R., and Ylitalo, G.M., 2010. Status Review of Hawaiian Insular False Killer Whales (*Pseudorca crassidens*) under the Endangered Species Act. U.S. Dep. Commer. NOAA Tech Memo., NOAA-TM-NMFS-PIFSC-22. 140 pp.
- Oleson, E.M., Baird, R.W., Martien, K.K., and Taylor, B.L., 2013. Island-associated stocks of odontocetes in the main Hawaiian Islands: A synthesis of available information to facilitate evaluation of stock structure. PIFSC Working Paper WP-13-003. 41 pp.

- Olson, P.A., Ensor, P., Olavarria, C., Bott, N., Constantine, R., Weir, J., Childerhouse, S., van der Linde, M., Schmitt, N., Miller, B.S. and Double, M.C., 2015. New Zealand Blue Whales: Residency, Morphology, and Feeding Behavior of a Little-Known Population. *Pacific Science*, 69(4), pp. 477-485.
- Perrin, W.F., Wursig, B., and Thewissen J.G.M. [eds.], 2009. Encyclopedia of marine mammals. *Academic Press*.
- Perryman, W.L., Au, D.W.K., Leatherwood, S., and Jefferson, T.A., 1994. Melon-headed whale *Peponocephala electra* Gray, 1846. In: Ridgway, S.H. and Harrison, R. [eds.]. Handbook of Marine Mammals, Volume 5. The first book of dolphins. *Academic Press*, London, U.K.
- Pitman, R.L. 1990. Pelagic distribution and biology of sea turtles in the eastern tropical Pacific. In: Richardson, T.H., Richardson, J.I., and Donnelly, M. [eds.]. Proc. of the Tenth Annual Workshop on Sea Turtle Biology and Conservation. U.S. Dep. of Comm., NOAA Tech. Memo. NMFS-SEFC-278. 286 pp.
- Pyle, R.L., and Pyle, P., 2009. The Birds of the Hawaiian Islands: Occurrence, History, Distribution, and Status. B.P. Bishop Museum, Honolulu, Hawaii. Version 1 (31 December 2009). Accessed from <http://hbs.bishopmuseum.org/birds/rlp-monograph>.
- Rankin, S. and Barlow, J., 2005. Source of the North Pacific “boing” sound attributed to minke whales. *Journal of the Acoustical Society of America*, 118(5), pp. 3346-3335
- Reeves, R.R., Leatherwood, S., Stone, G.S., and Eldredge, L.G., 1999. Marine Mammals in the Area Served by the South Pacific Regional Environment Program. Report of the South Pacific Regional Environment Program (P.O. Box 240, Apia, Samoa). 48 pp.
- Reeves, R.R., Leatherwood, S., and Baird, R.W., 2009. Evidence of a possible decline since 1989 in false killer whales (*Pseudorca crassidens*) around the main Hawaiian Islands. *Pacific Science*, 63(2), pp. 253–261.
- Rice, D.W., 1960. Distribution of the bottle-nosed dolphin in the leeward Hawaiian Islands. *Journal of Mammalogy*, 41(3), pp. 407-408.
- Rice, D. W., and A. A. Wolman. 1984. Humpback whale census in Hawaiian waters—February 1977. In: Norris, K.S., and Reeves, R.R. (eds.). Report on a workshop on problems related to humpback whales (*Megaptera novaeangliae*) in Hawaii. Final report to the Marine Mammal Commission, U.S. Department of Commerce, NTIS PB-280-794.
- Ross, G.J.B. and Leatherwood, S., 1994. Pygmy killer whale *Feresa attenuata* (Gray, 1874). Handbook of Marine Mammals, Volume 5, pp. 387-404.
- Rowntree, V., Darling, J., Silber, G., and M. Ferrari, M., 1980. Rare sighting of a right whale (*Eubalaena glacialis*) in Hawaii. *Canadian Journal of Zoology*, 58, pp. 308-312.

- Sanches, J.G., 1991. Catálogo dos principais peixes marinhos da República de Guiné-Bissau. Publicações avulsas do I.N.I.P. No. 16. 429 pp. as cited in Froese, R. and D. Pauly, Editors. 2000. FishBase 2000: concepts, design and data sources. ICLARM, Los Baños, Laguna, Philippines. 344 pp.
- Schorr, G.S., Baird, R.W., Hanson, M.B., Webster, D.L., McSweeney, D.J., and Andrews, R.D., 2009. Movements of satellite-tagged Blainville's beaked whales off the island of Hawai'i. *Endangered Species Research*, 10, pp. 203-213.
- Schulze-Haugen, M. and Kohler, N.E. [eds.], 2003. Guide to Sharks, Tunas, & Billfishes of the U.S. Atlantic and Gulf of Mexico. RI Sea Grant/National Marine Fisheries Service.
- Shallenberger, E.W., 1981. The status of Hawaiian cetaceans. Marine Mammal Commission Report No. MMC-77/23 (NTIS PB82-109398).
- Stafford, K.M., Nieuwirth, S.L., and Fox, C.G., 2001. Geographic and seasonal variation of blue whale calls in the North Pacific. *Journal of Cetacean Research and Management*, 3, pp. 65-76.
- Strasburg, D.W., 1958. Distribution, abundance, and habits of pelagic sharks in the central Pacific Ocean. *Fishery Bulletin*, 138(58), pp. 335-361.
- Thompson, P.O., and Friedl, W.A., 1982. A long-term study of low frequency sounds from several species of whales off Oahu, Hawaii. *Cetology*, 45, pp. 1-19.
- Tuato'o-Bartley, N., Morrell, T.E., and Craig, P., 1993. Status of sea turtles in American Samoa in 1991. *Pacific Science*, 47(3), pp. 215-221.
- Utzurum, R., 2002. Sea turtle conservation in American Samoa. In: Kinan, I. [ed.]. Proceedings of the western Pacific sea turtle cooperative research and management workshop.
- Veron, J.E.N., 2014. Results of an update of the Corals of the World Information Base for the Listing Determination of 66 Coral Species under the Endangered Species Act. Report to the Western Pacific Regional Fishery Management Council, Honolulu, Hawaii.
- Wolman, A.A. and Jurasz, C.M., 1977. Humpback whales in Hawaii: Vessel Census, 1976. *Marine Fisheries Review*, 39(7), pp. 1-5.