

2.1 CORAL REEF FISH ECOSYSTEM PARAMETERS

2.1.1 Regional Reef Fish Biomass and Habitat Condition

Description: ‘Reef fish biomass’ is mean biomass of reef fishes per unit area derived from visual survey data between 2010 and 2020. Hard Coral cover is mean cover derived from visual estimates by divers of sites where reef fish surveys occurred. No new survey occurred in 2020 and the numbers presented here are identical to last year’s report.

Rationale: Reef fish biomass has been widely used as an indicator of relative ecosystem status and has repeatedly been shown to be sensitive to changes in fishing pressure, habitat quality, and oceanographic regime. Hard coral cover is an indicator of relative status of the organisms that build coral reef habitat and has been shown to be sensitive to changes in oceanographic regime, and a range of direct and indirect anthropogenic impacts. Most fundamentally, cover of hard corals has been increasingly impacted by temperature stress as a result of global heating.

Data Category: Fishery-independent

Timeframe: Triennial

Jurisdiction: American Samoa, Guam, Commonwealth of the Northern Mariana Islands (CNMI), Main Hawaiian Islands (MHI), Northwestern Hawaiian Islands (NWHI), and Pacific Remote Island Areas (PRIAs)

Spatial Scale: Regional

Data Source: Data used to generate cover and biomass estimates come from visual surveys conducted by the National Marine Fisheries Service (NMFS) Pacific Island Fisheries Science Center (PIFSC) Ecosystem Sciences Division (ESD) and their partners as part of the Pacific Reef Assessment and Monitoring Program ([RAMP](#)). Survey methods are described in detail in Ayotte et al. (2015). In brief, they involve teams of divers conducting stationary point count cylinder (SPC) surveys within a target domain of < 30 meter hard-bottom habitat at each island, stratified by depth zone and, for larger islands, by section of coastline. For consistency among islands, only data from forereef habitats are used. At each SPC, divers record the number, size, and species of all fishes within or passing through paired 15 meter-diameter cylinders over the course of a standard count procedure.

Fish sizes and abundance are converted to biomass using standard length-to-weight conversion parameters, taken largely from [FishBase](#) and converted to biomass per unit area by dividing by the area sampled per survey. Site-level data were pooled into island-scale values by first calculating mean and variance within strata, and then calculating weighted island-scale mean and variance using the formulas given in Smith et al. (2011) with strata weighted by their respective sizes.




Figure 1. Mean coral cover (%) per U.S. Pacific island averaged over the years 2010-2020 by latitude

DRAFT




Figure 2. Mean fish biomass (g/m² ± standard error) of functional, taxonomic and trophic groups by U.S. Pacific reef area from the years 2010-2020 by latitude. The group Serranidae excludes planktivorous members of that family – i.e. anthias, which can be hyper-abundant in some regions. Similarly, the bumphead parrotfish, *Bolbometopon muricatum*, has been excluded from the corallivore group – as high biomass of that species at Wake overwhelms corallivore biomass at all other locations. The group ‘MI Feeder’ consists of fishes that primarily feed on mobile invertebrates

2.1.2 Main Hawaiian Islands Reef Fish Biomass and Habitat Condition

Description: ‘Reef fish biomass’ is mean biomass of reef fishes per unit area derived from visual survey data between 2010 and 2020. Hard Coral cover is mean cover derived from visual estimates by divers of sites where reef fish surveys occurred. No new survey occurred in 2020 and the numbers presented here are identical to last year’s report.

Rationale: Reef fish biomass has been widely used as an indicator of relative ecosystem status and has repeatedly been shown to be sensitive to changes in fishing pressure, habitat quality, and oceanographic regime. Hard coral cover is an indicator of relative status of the organisms that build coral reef habitat and has been shown to be sensitive to changes in oceanographic regime, and a range of direct and indirect anthropogenic impacts. Most fundamentally, cover of hard corals has been increasingly impacted by temperature stress as a result of global heating.

Data Category: Fishery-independent

Timeframe: Triennial

Jurisdiction: MHI

Spatial Scale: Island

Data Source: Data used to generate biomass and cover estimates comes from visual surveys conducted by NOAA PIFSC ESD and their partners, as part of the Pacific RAMP. Survey methods and sampling design, and methods to generate reef fish biomass are described in Section 2.1.1.




Figure 3. Mean coral cover (%) per island averaged over the years 2010-2020 by latitude with MHI mean estimates plotted for reference (red line)

DRAFT




Figure 4. Mean fish biomass ($\text{g}/\text{m}^2 \pm$ standard error) of MHI functional, taxonomic and trophic groups from the years 2010-2020 by island. The group Serranidae excludes planktivorous members of that family – i.e. anthias, which can be hyper-abundant in some regions. The group ‘MI Feeder’ consists of fishes that primarily feed on mobile invertebrates; with MHI mean estimates plotted for reference (red line)

2.1.3 Northwestern Hawaiian Islands Reef Fish Biomass and Habitat Condition

Description: ‘Reef fish biomass’ is mean biomass of reef fishes per unit area derived from visual survey data between 2010 and 2020. Hard Coral cover is mean cover derived from visual estimates by divers of sites where reef fish surveys occurred. No new survey occurred in 2020 and the numbers presented here are identical to last year’s report.

Rationale: Reef fish biomass has been widely used as an indicator of relative ecosystem status and has repeatedly been shown to be sensitive to changes in fishing pressure, habitat quality, and oceanographic regime. Hard coral cover is an indicator of relative status of the organisms that build coral reef habitat and has been shown to be sensitive to changes in oceanographic regime, and a range of direct and indirect anthropogenic impacts. Most fundamentally, cover of hard corals has been increasingly impacted by temperature stress as a result of global heating.

Data Category: Fishery-independent

Timeframe: Triennial

Jurisdiction: NWHI

Spatial Scale: Island

Data Source: Data used to generate biomass and cover estimates comes from visual surveys conducted by NOAA PIFSC ESD and their partners, as part of the Pacific RAMP. Survey methods and sampling design, and methods to generate reef fish biomass are described above (Section 2.1.1).




Figure 5. Mean coral cover (%) per island averaged over the years 2010-2020 by latitude with NWHI mean estimates plotted for reference (red line)




Figure 6. Mean fish biomass ($\text{g}/\text{m}^2 \pm$ standard error) of NWHI functional, taxonomic and trophic groups from the years 2010-2020 by island. The group Serranidae excludes planktivorous members of that family – i.e. anthias, which can be hyper-abundant in some regions. The group ‘MI Feeder’ consists of fishes that primarily feed on mobile invertebrates; with NWHI mean estimates plotted for reference (red line)